

A close-up photograph of a hand in a white shirt cuff striking a matchstick. The matchstick is lit, with a bright yellow flame and wisps of white smoke. To the left of the lit matchstick, there is a row of seven unlit matchsticks standing upright on a dark, textured wooden surface. The background is a blurred wooden surface.

Causal Inference – Theory and Applications

Dr. Matthias Uflacker, Johannes Huegle, Christopher Schmidt

April 24, 2018

- **Jupyter Notebook „Causal Inference in Application“**
- **Recap Causal Inference in a Nutshell**
- **Introduction to Structural Causal Models**
 1. Preliminaries
 2. Structural Causal Models
 3. (Local) Markov Condition
 4. Factorization
 5. Global Markov Condition
 6. Functional Model and Markov conditions
 7. Faithfulness
 8. Constraint-based Causal Inference
 9. Markov Equivalence Class
 10. Summary
 11. Excursion: Maximal Ancestral Graphs

**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

Slide 2

The image shows a screenshot of the RStudio interface. The main editor window displays R code for loading and analyzing the 'diamonds' dataset. The code includes loading 'ggplot2', sourcing a plot format file, viewing the dataset, and creating a plot. The plot title is 'Diamond Pricing'. The right-hand pane shows the 'Workspace' and 'History' tabs, with the 'Data' section displaying the 'diamonds' dataset (53940 observations) and the 'Functions' section showing the 'format.plot' function. The bottom of the plot area is partially obscured by a red overlay.

```
1 library(ggplot2)
2 source("plots/formatPlot.R")
3
4 view(diamonds)
5 summary(diamonds)
6
7 summary(diamonds$price)
8 aveSize <- round(mean(diamonds$carat), 4)
9 aveClarity <- round(mean(diamonds$clarity), 4)
10
11 plot(diamonds, aes(carat, price),
12      data=data=diamonds, color=clarity,
13      xlab="Carat", ylab="Price",
14      main="Diamond Pricing")
15
```

Workspace History
Load Save Import Dataset Clear All
Data
diamonds 53940 obs. of 10 variables
Values
aveSize 0.7979
clarity character [8]
p ggplot [8]
Functions
format.plot(plot, size)

Files Plots Packages Help
Zoom Export Clear All

Diamond Pricing

Jupyter Notebook

"Causal Inference in Application"

Jupyter Notebook

Causal Inference in Application

Causal Inference - Theory and Applications

In our lecture [Causal Inference - Theory and Applications](#), we look at the mathematical concepts that build the basis of causal inference.



Causal Inference in Application

We now look how these concepts are applied on observational data to derive causal relationships and how to use the do-operator to receive an estimation of the causal effect. In order to give you an overview on the related procedure, this notebook gives a step by step approach in the context of a simple cooling house example.

Table of Contents

1. [Introduction to R](#)
 - A. [Getting Started](#)
 - B. [Some Examples](#)
2. [Use Case](#)
 - A. [Description](#)

Causal Inference - Theory and Applications

Uflacker, Huegle,
Schmidt

Slide 4

Jupyter Notebook


Access Information

System

Link will be provided via email once we have the list of participants!

Procedure

1. Login via LDAP (standard HPI credentials)
2. Use folder Causal Inference – Theory and Applications
3. We provide a Master Notebook
Please use as a read only resource
Copy relevant information into your local workspace
4. Your local workspace either in your home directory or as a separate folder in our courses' folder
5. Let us know if you require new packages



**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

Slide 5

A close-up photograph of a hand in a dark suit jacket and white shirt cuff, striking a matchstick. The match is lit, with a bright yellow flame and wisps of white smoke. To the left of the lit match are seven other unlit matchsticks, all standing upright on a dark, textured wooden surface. The background is a blurred wooden wall with a prominent grain pattern. A semi-transparent red banner is overlaid at the bottom of the image, containing the title text.


Causal Inference in a Nutshell

Causal Inference in a Nutshell

Recap: The Concept

Traditional Statistical Inference Paradigm

Paradigm of Structural Causal Models



E.g., what is the sailors' probability of recovery when **we see** a treatment with lemons?

$$Q(P) = P(\text{recovery}|\text{lemons})$$


E.g., what is the sailors' probability of recovery if **we do** treat them with lemons?

$$Q(G) = P(\text{recovery}|\text{do}(\text{lemons}))$$


Causal Inference
- Theory and Applications

Uflacker, Huegle,
Schmidt

Slide 7



Introduction to Structural Causal Models



Introduction to Causal Graphical Models

Content

1. Preliminaries
2. Structural Causal Models
3. (Local) Markov Condition
4. Factorization
5. Global Markov Condition
6. Functional Model and Markov conditions
7. Faithfulness
8. Constraint-based Causal Inference
9. Markov Equivalence Class
10. Summary
11. Excursion: Maximal Ancestral Graphs

**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

1. Preliminaries

Notation

- A, B events
- X, Y, Z random variables
- x value of random variable

- Pr probability measure
- P_X probability distribution of X
- p density
- p_x or $p(X)$ density of P_X
- $p(x)$ density of P_X evaluated at the point x

- $X \perp Y$ independence of X and Y
- $X \perp Y \mid Z$ conditional independence of X and Y given Z

1. Preliminaries

Independence of Events

- Two events A and B are called *independent* if

$$\Pr(A \cap B) = \Pr(A) \cdot \Pr(B),$$

or - rewritten in *conditional probabilities* - if

$$\Pr(A) = \frac{\Pr(A \cap B)}{\Pr(B)} = \Pr(A|B),$$

$$\Pr(B) = \frac{\Pr(A \cap B)}{\Pr(A)} = \Pr(B|A).$$

- A_1, \dots, A_n are called (*mutually*) *independent* if for every subset $S \subset \{1, \dots, n\}$ we have

$$\Pr\left(\bigcap_{i \in S} A_i\right) = \prod_{i \in S} \Pr(A_i).$$

- Note:**

for $n \geq 3$, pairwise independence $\Pr(A_i \cap A_j) = \Pr(A_i) \cdot \Pr(A_j)$ for all i, j does not imply (mutual) independence.

1. Preliminaries

Independence of Random Variables

- Two real-valued random variables X and Y are called *independent*,

$$X \perp Y,$$

if for every $x, y \in \mathbb{R}$, the events $\{X \leq x\}$ and $\{Y \leq y\}$ are independent,

Or, in terms of densities: for all x, y ,

$$p(x, y) = p(x)p(y).$$

- Note:**

If $X \perp Y$, then $E[XY] = E[X]E[Y]$, and $cov(X, Y) = E[XY] - E[X]E[Y] = 0$.

The converse is not true: If, $cov(X, Y) = 0$, then $X \perp Y$.

No correlation does not imply dependence

However, we have, for large \mathcal{F} : $(\forall f, g \in \mathcal{F}: cov(f(X), g(Y)) = 0)$, then $X \perp Y$.

Causal Inference
- Theory and
Applications

Uflacker, Huegle,
Schmidt

Slide 12

1. Preliminaries

Conditional Independence of Random Variables

- Two real-valued random variables X and Y are called *conditionally independent* given Z ,

$$X \perp Y \mid Z \text{ or } (X \perp Y \mid Z)_p$$

if

$$p(x, y|z) = p(x|z)p(y|z)$$

For all x, y and for all z s.t. $p(z) > 0$.

- Note:**

It is possible to find X, Y which are conditionally independent given a variable Z but unconditionally dependent, and vice versa.


2. Structural Causal Models

Definition (Pearl)

- Directed Acyclic Graph (DAG) $G = (V, E)$
 - *Vertices* V_1, \dots, V_n
 - *Directed edges* $E = (V_i, V_j)$, i.e., $V_i \rightarrow V_j$,
 - *No cycles*
- Use kinship terminology, e.g., for path $V_i \rightarrow V_j \rightarrow V_k$
 - $V_i = Pa(V_j)$ *parent* of V_j
 - $\{V_i, V_j\} = Ang(V_k)$ *ancestors* of V_k
 - $\{V_j, V_k\} = Des(V_i)$ *descendants* of V_i
- Directed Edges encode *direct causes* via
 - $V_j = f_j(Pa(V_j), N_j)$ with independent noise N_1, \dots, N_n

➔ This forms the Causal Graphical Model

Cooling House Example:



- $V_1 = N(0,1)$
- $V_2 = N(0,1)$
- $V_3 = 3 V_2 + N(0,1)$
- $V_4 = 4 V_1 + 5 V_2 + 0.7 V_3 + N(0,1)$
- $V_5 = V_4 + N(0,1)$
- $V_6 = 1.2 V_4 + N(0,1)$


Causal Inference - Theory and Applications

Uflacker, Huegle,
Schmidt

2. Structural Causal Models

Connecting G and P

- Basic Assumption: *Causal Sufficiency*
 - All relevant variables are included in the DAG G



$$(X \perp Y|Z)_G \Rightarrow (X \perp Y|Z)_P$$

- Key Postulate: *(Local) Markov Condition*
- Essential mathematical concept: *d-separation*
(describes the conditional independences required by a causal DAG)

Causal Inference
- Theory and
Applications

Uflacker, Huegle,
Schmidt

Slide 15


3. (Local) Markov Condition Theorem

(Local) Markov Condition:

V_j statistically independent of nondescendants, given parents $Pa(V_j)$, i.e.,

$$V_j \perp V_{V/Des(V_j)} | Pa(V_j).$$

- I.e., every information exchange with its nondescendants involves its parents
- Example:



- $V_6 \perp \{V_1, V_2, V_3\} | V_4$
- $V_5 \perp \{V_1, V_2, V_3\} | V_4$

Causal Inference
- Theory and
Applications

Uflacker, Huegle,
Schmidt

Slide 16

3. (Local) Markov Condition Supplement (Lauritzen 1996)

- Assume V_n has no descendants, then $ND_n = \{V_1, \dots, V_{n-1}\}$.

- Thus the local Markov condition implies

$$V_n \perp \{V_1, \dots, V_{n-1}\} | Pa(V_n).$$

- Hence, the general decomposition

$$p(v_1, \dots, v_n) = p(v_n | v_1, \dots, v_{n-1}) p(v_1, \dots, v_{n-1})$$

becomes

$$p(v_1, \dots, v_n) = p(v_n | Pa(v_n)) p(v_1, \dots, v_{n-1}).$$

- Induction over n yields to

$$p(v_1, \dots, v_n) = \prod_{i=1}^n p(v_i | Pa(v_i)).$$

- I.e., the graph shows us how to factor the joint distribution P_V .


4. Factorization

Definition

Factorization:

$$p(v_1, \dots, v_n) = \prod_{i=1}^n p(v_i | Pa(v_i)).$$

- I.e., conditionals as causal mechanisms generating statistical dependence
- Example:



$$\begin{aligned} p(V) &= p(v_1, \dots, v_n) \\ &= p(v_1) \cdot p(v_2) \\ &\quad \cdot p(v_3 | v_2) \cdot p(v_4 | v_1, v_2, v_3) \\ &\quad \cdot p(v_5 | v_4) \cdot p(v_6 | v_4) \\ &= \prod_{i=1}^n p(v_i | Pa(v_i)) \end{aligned}$$

5. Global Markov Condition

D-Separation (Pearl 1988)

- *Path* = sequence of pairwise distinct vertices where consecutive ones are adjacent
- A path q is said to be *blocked* by a set S if
 - q contains a *chain* $V_i \rightarrow V_j \rightarrow V_k$ or a *fork* $V_i \leftarrow V_j \rightarrow V_k$ such that the middle node is in S , or
 - q contains a *collider* $V_i \rightarrow V_j \leftarrow V_k$ such that the middle node is not in S and such that no descendant of V_j is in S .

D-separation:

S is said to **d-separate** X and Y in the DAG G , i.e.,

$$(X \perp Y | S)_G,$$

if S blocks every path from a vertex in X to a vertex in Y .

Causal Inference
- Theory and
Applications


Uflacker, Huegle,
Schmidt

Slide 19

5. Global Markov Condition

Examples of d-Separation

- Example:




- The path from V_1 to V_6 is blocked by V_4 .
- V_1 and V_6 are d-separated by V_4 .
- The path $V_2 \rightarrow V_3 \rightarrow V_4 \rightarrow V_6$ is blocked by V_3 or V_4 or both.
- But: V_2 and V_6 are d-separated only by V_4 or $\{V_3, V_4\}$.
- V_1 and V_2 are not blocked by V_4 .

5. Global Markov Condition Theorem

Global Markov Condition:

For all disjoint subsets of vertices X, Y and Z we have that
 X, Y d-separated by $Z \Rightarrow (X \perp Y | Z)_P$.

- I.e., we have $(X \perp Y | Z)_G \Rightarrow (X \perp Y | Z)_P$



Causal Inference
- Theory and
Applications

Uflacker, Huegle,
Schmidt

Slide 21

6. Functional Model and Markov Conditions Theorem (Lauritzen 1996, Pearl 2000)

Theorem:

The following are equivalent:

- Existence of a *functional causal model* G ;
- *Local Causal Markov condition*: V_j statistically independent of nondescendants, given parents
(i.e.: every information exchange with its nondescendants involves its parents)
- *Global Causal Markov condition*: d-separation
(characterizes the set of independences implied by local Markov condition)
- *Factorization*: $p(v_1, \dots, v_n) = \prod_{i=1}^n p(v_i | Pa(v_i))$.

(subject to technical conditions)

$$\text{i.e., } (X \perp Y | Z)_G \Rightarrow (X \perp Y | Z)_P$$

**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

Slide 22

7. Causal Faithfulness


The key-postulate

Causal Faithfulness:

p is called faithful relative to G if only those independencies hold true that are implied by the Markov condition, i.e.,

$$(X \perp Y | Z)_G \Leftarrow (X \perp Y | Z)_P$$

- I.e., we assume that any population P produced by this causal graph G has the independence relations obtained by applying d-separation to it
- Seems like a hefty assumption, but it really isn't: It assumes that whatever independencies occur in it arise not from incredible coincidence but rather from structure, i.e., data generating model G .
- Hence:



Causal Inference
- Theory and
Applications

Uflacker, Huegle,
Schmidt

Slide 23

8. Constraint-based Causal Inference

Concept (Spirtes, Glymour, Scheines and Pearl)

■ Assumptions:

- Causal Sufficiency
- Global Markov Condition
- Causal Faithfulness

■ Causal Structure Learning:

- Accept only those DAG's G as causal hypothesis for which
$$(X \perp Y | Z)_G \Leftrightarrow (X \perp Y | Z)_P.$$
- Defines the basis of constraint-based causal structure learning
- Identifies causal DAG up to Markov equivalence class (DAGs that imply the same conditional independencies)

**Causal Inference
- Theory and
Applications**


Uflacker, Huegle,
Schmidt

9. Markov Equivalence Class Theorem (Verma and Pearl)

Theorem:

Two DAGs are Markov equivalent if and only if they have the same skeleton and the same v -structures


- *Skeleton:*
corresponding undirected graph
- *v -structure:*
substructure $X \rightarrow Y \leftarrow Z$ with no edges between X and Z .




9. Markov Equivalence Class

Examples

- Same skeleton, no v -structure
- Same skeleton, same v -structure at W



$$X \perp Z \mid Y$$



10. Summary

Causal Structural Models

- Causal Structures formalized by DAG (directed acyclic graph) G with random variables V_1, \dots, V_n as vertices.
- Causal Sufficiency, Causal Faithfulness and Markov Condition imply
$$(X \perp Y | Z)_G \Leftrightarrow (X \perp Y | Z)_P.$$
- Local Markov Condition states that the density $p(v_1, \dots, v_n)$ then factorizes into
$$p(v_1, \dots, v_n) = \prod_{i=1}^n p(v_i | Pa(v_i)).$$
- Causal conditional $p(v_j | Pa(v_j))$ represent causal mechanisms.

11. Excursion: Maximal Ancestral Graphs

Motivating Example

- Suppose, we are given the following list of conditional independencies among X, Y, Z and W :

- $X \perp\!\!\!\perp Z$,
- $Y \perp\!\!\!\perp W$,
- $X \perp\!\!\!\perp W$.
- $X \not\perp\!\!\!\perp Y$,
- $Y \not\perp\!\!\!\perp Z$,
- $Z \not\perp\!\!\!\perp W$.

- Which DAG could have generated these, and only these, independencies and dependencies?
- The pattern of dependencies must be:

$$X \text{ --- } Y \text{ --- } Z \text{ --- } W$$

- And there must be the following colliders:

$$X \longrightarrow Y \longleftarrow Z$$


$$Y \longrightarrow Z \longleftarrow W$$

- There is no orientation of Y – Z that is consistent with the independencies.

11. Excursion: Maximal Ancestral Graphs

DAG Models and Marginalization

- Let's include an additional variable U :



- This DAG model generates a probability distribution $P_{\{X,Y,Z,W,U\}}$ in which:

- $X \perp\!\!\!\perp Z$,
- $X \not\perp\!\!\!\perp Y$,
- $Y \perp\!\!\!\perp W$,
- $Y \not\perp\!\!\!\perp Z$,
- $X \perp\!\!\!\perp W$.
- $Z \not\perp\!\!\!\perp W$.

- The marginal distribution $P_{\{X,Y,Z,W\}} = P_{\{X,Y,Z,W,U\}} du$ must adhere the same independencies. But: this marginal distribution cannot be faithfully generated by any DAG.


➔ DAG models are not closed under marginalization!

11. Excursion: Maximal Ancestral Graphs

Ancestral Graphs (informally)

- Ancestral Graph (AG)**


is a graph containing both directed and bi-directed edges, where the bi-directed edges stand for *latent variables*, e.g.,



- m-Separation**

If S m-separates X and Y in an ancestral graph M , then $X \perp Y \mid S$ in every density p that factorizes according to any DAG G that is represented by the AG M .

- Example**



**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

11. Excursion: Maximal Ancestral Graphs

DAGs vs. AGs

■ Advantages of AGs

- AGs can faithfully represent more probability distributions than DAGs.
- AG models are closed under marginalization.
- AGs can (implicitly) represent unobserved variables, which exist in many (possibly almost all) applications.

■ Disadvantages of AGs

- Parameterization is difficult in the general case.
- Markov equivalence is difficult.

**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

Slide 31

Literature

- Pearl, J. (2009). *Causal inference in statistics: An overview*. Statistics Surveys, 3: 96-146.
- Pearl, J. (2009). *Causality: Models, Reasoning, and Inference*. Cambridge University Press.
- Spirtes, P., Glymour, C., and Scheines, R. (2000). Causation, Prediction, and Search. The MIT Press.

**Causal Inference
- Theory and
Applications**

Uflacker, Huegle,
Schmidt

Slide 32

Thank you
for your attention!