

1 Rainbow Vertex Coloring Bipartite Graphs and 2 Chordal Graphs

3 **Pinar Heggernes**

4 Department of Informatics, University of Bergen, Norway

5 pinar.heggernes@uib.no

6 **Davis Issac**

7 Max Planck Institute for Informatics, Saarland Informatics Campus, Germany

8 dissac@mpi-inf.mpg.de

9 **Juho Lauri**

10 Nokia Bell Labs, Dublin, Ireland

11 juho.lauri@nokia-bell-labs.com

12 **Paloma T. Lima**

13 Department of Informatics, University of Bergen, Norway

14 paloma.lima@uib.no

15 **Erik Jan van Leeuwen**

16 Department of Information and Computing Sciences, Utrecht University, The Netherlands

17 e.j.vanleeuwen@uu.nl

18 — Abstract —

19 Given a graph with colors on its vertices, a path is called a rainbow vertex path if all its internal
20 vertices have distinct colors. We say that the graph is rainbow vertex-connected if there is
21 a rainbow vertex path between every pair of its vertices. We study the problem of deciding
22 whether the vertices of a given graph can be colored with at most k colors so that the graph
23 becomes rainbow vertex-connected. Although edge-colorings have been studied extensively under
24 similar constraints, there are significantly fewer results on the vertex variant that we consider. In
25 particular, its complexity on structured graph classes was explicitly posed as an open question.

26 We show that the problem remains NP-complete even on bipartite apex graphs and on split
27 graphs. The former can be seen as a first step in the direction of studying the complexity of
28 rainbow coloring on sparse graphs, an open problem which has attracted attention but limited
29 progress. We also give hardness of approximation results for both bipartite and split graphs. To
30 complement the negative results, we show that bipartite permutation graphs, interval graphs,
31 and block graphs can be rainbow vertex-connected optimally in polynomial time.

32 **2012 ACM Subject Classification** G.2.2 Graph Theory

33 **Keywords and phrases** Rainbow coloring, graph classes, polynomial-time algorithms, approxi-
34 mation algorithms

35 **Digital Object Identifier** 10.4230/LIPIcs.MFCS.2018.83

36 **1** Introduction

37 Graph coloring and graph connectivity are two of the most famous topics in graph algorithms.
38 Many different types of colorings and connectivity measures have been considered throughout
39 time. The concept of rainbow coloring brings these two extensively studied topics together,
40 and it was first defined a decade ago by Chartrand et al. [8] using edge-colorings. Let G be a
41 connected, edge-colored graph. A *rainbow path* in G is a path all of whose edges are colored



© Pinar Heggernes, Davis Issac, Juho Lauri, Paloma T. Lima, Erik Jan van Leeuwen;
licensed under Creative Commons License CC-BY

43rd International Symposium on Mathematical Foundations of Computer Science (MFCS 2018).

Editors: Igor Potapov, Paul Spirakis, and James Worrell; Article No. 83; pp. 83:1–83:13

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

with distinct colors, and G is *rainbow-connected* if there is a rainbow path between every pair of its vertices. The resulting computational problem RAINBOW COLORING (RC) takes as input a connected (uncolored) graph G and an integer k , and the task is to decide whether the edges of G can be colored with at most k colors so that G is rainbow-connected. This problem has various applications in telecommunications, data transfer, and encryption [25, 4, 11] and has been studied rather thoroughly from both graph-theoretic and complexity-theoretic viewpoints (see related work below and the surveys [19, 25]).

The intense interest in RAINBOW COLORING led Krivelevich and Yuster [18] to define a natural variant on *vertex-colored* graphs. Here, a path in a vertex-colored graph H is a *rainbow vertex path* if all its *internal* vertices have distinct colors. We say that H is *rainbow vertex-connected* if there is a rainbow vertex path between every pair of its vertices. Similarly to the edge variant, RAINBOW VERTEX COLORING (RVC) is the decision problem in which we are given a connected (uncolored) graph H and an integer k , and the task is to decide whether the vertices of H can be colored with at most k colors such that H is rainbow vertex-connected. The *rainbow vertex connection number* of G , denoted by $\mathbf{rvc}(G)$, is the minimum k such that G has a rainbow vertex coloring with k colors. RVC is NP-complete for every $k \geq 2$ [10, 9], and remains NP-complete for $k = 3$ for bipartite graphs [23]. In addition, it is NP-hard to approximate $\mathbf{rvc}(G)$ within a factor of $2 - \varepsilon$ unless $\mathsf{P} \neq \mathsf{NP}$, for any $\varepsilon > 0$ [13]. It is also known that RVC is linear-time solvable on planar graphs for every fixed k [19]. Finally, assuming the Exponential Time Hypothesis, there is no algorithm for solving RVC in time $2^{o(n^{3/2})}$ for any $k \geq 2$ [19].

A stronger variant of rainbow vertex-colorings was introduced by Li et al. [24]. A vertex-colored graph H is *strongly rainbow vertex-connected* if between every pair of vertices of H , there is a *shortest* path that is also a rainbow vertex path. The STRONG RAINBOW VERTEX COLORING (SRVC) problem takes as input a connected (uncolored) graph H and an integer k , and the task is to decide whether the vertices of H can be colored such that H is strongly rainbow vertex-connected. This definition is the vertex variant of the STRONG RAINBOW COLORING problem, which was also broadly studied (see related work below and the surveys [19, 25]). The *strong rainbow vertex connection number* of G , denoted by $\mathbf{srvc}(G)$, is the minimum k such that G has a strong rainbow vertex coloring with k colors. SRVC is NP-complete for every $k \geq 2$ [12] and linear-time solvable on planar graphs for every fixed k [19]. In addition, it is NP-hard to approximate $\mathbf{srvc}(G)$ within a factor of $n^{1/2-\varepsilon}$ unless $\mathsf{P} \neq \mathsf{NP}$, for any $\varepsilon > 0$ [13].

While RC has been widely studied in more than 300 published papers, we are unaware of any further complexity results on RVC and SRVC than those mentioned previously. In particular, the complexity of RVC and SRVC on structured graph classes is mostly open. This led Lauri [19, Open problem 6.6] to explicitly ask the following:

For what restricted graph classes do RVC and SRVC remain NP-complete?

Our Results In this paper, we make significant progress towards addressing this open problem. In particular, we study bipartite graphs and chordal graphs, and some of their subclasses, and give hardness results and polynomial-time algorithms for RVC and SRVC. Our main result is a hardness result for bipartite apex graphs:

► **Theorem 1.** *Let G be a bipartite apex graph of diameter 4. It is NP-complete to decide both whether $\mathbf{rvc}(G) \leq 4$ and whether $\mathbf{srvc}(G) \leq 4$. Moreover, it is NP-hard to approximate $\mathbf{rvc}(G)$ and $\mathbf{srvc}(G)$ within a factor of $5/4 - \varepsilon$, for every $\varepsilon > 0$.*

This result is particularly interesting since no hardness result was known on a sparse graph class (like apex graphs) for any of the variants of rainbow coloring. Moreover, this result

89 can be considered tight in conjunction with the known result that RVC and SRVC are
 90 linear-time solvable on planar graphs for every fixed number of colors k [19]. Finally, we
 91 observe (like Li et al. [23]) that $\mathbf{rvc}(G)$ and $\mathbf{srvc}(G)$ can be computed in linear time if G is
 92 a bipartite graph of diameter 3, providing further evidence that this result is tight.

93 For general bipartite graphs and for split graphs (a well-known subclass of chordal graphs),
 94 we exhibit stronger hardness results:

95 ► **Theorem 2.** *Let G be a bipartite graph of diameter 4. It is NP-complete to decide both*
 96 *whether $\mathbf{rvc}(G) \leq k$ and whether $\mathbf{srvc}(G) \leq k$, for every $k \geq 3$. Moreover, it is NP-hard to*
 97 *approximate both $\mathbf{rvc}(G)$ and $\mathbf{srvc}(G)$ within a factor of $n^{1/3-\varepsilon}$, for every $\varepsilon > 0$.*

98 We remark that, previously, it was only known that deciding whether $\mathbf{rvc}(G) \leq 3$ for bipartite
 99 graphs G is NP-complete by the result of [23]. Our construction, however, is conceptually
 100 simpler, gives hardness for every $k \geq 3$, and is easily extended to the strong variant. Moreover,
 101 for RVC on general graphs, this result implies a considerable improvement over the previous
 102 result of Eiben et al. [13] which only excluded a polynomial-time approximation with a factor
 103 of less than 2 assuming $P \neq NP$.

104 ► **Theorem 3.** *Let G be a split graph of diameter 3. It is NP-complete to decide both*
 105 *whether $\mathbf{rvc}(G) \leq k$ and whether $\mathbf{srvc}(G) \leq k$, for every $k \geq 2$. Moreover, it is NP-hard to*
 106 *approximate both $\mathbf{rvc}(G)$ and $\mathbf{srvc}(G)$ within a factor of $n^{1/3-\varepsilon}$, for every $\varepsilon > 0$.*

107 To the best of our knowledge, our results for split graphs give the first non-trivial graph class
 108 besides diameter-two graphs for which the complexity of the edge and the vertex variant
 109 differ (see e.g. [19, Table 4.2] but note that it contains a typo erroneously claiming that
 110 RVC can be solved in polynomial-time for split graphs). In particular, RC can be solved in
 111 polynomial time on split graphs when $k \geq 4$ [5, 7]. Moreover, we observe that $\mathbf{rvc}(G)$ and
 112 $\mathbf{srvc}(G)$ can be computed in linear time if G is a graph of diameter 2, providing evidence
 113 that this result is tight.

114 To contrast our hardness results, we show that both problems can be solved in polynomial
 115 time on several other subclasses of bipartite graphs and chordal graphs.

116 ► **Theorem 4.** *If G is a bipartite permutation graph, a block graph, or a unit interval graph,*
 117 *then $\mathbf{rvc}(G)$ and $\mathbf{srvc}(G)$ can be computed in linear time. If G is an interval graph, then*
 118 *$\mathbf{rvc}(G)$ can be computed in linear time.*

119 Combined, these results paint a much clearer picture of the complexity landscape of RVC
 120 and SRVC than was possible previously.

121 **Related Work** We briefly survey the known work for the edge variants of rainbow coloring;
 122 we refer to [19, 25] for more detailed surveys. RC is NP-complete for every $k \geq 2$ [4, 2, 22],
 123 even on chordal graphs [5]. On split graphs, RC is NP-complete when $k \in \{2, 3\}$, but solvable
 124 in polynomial time otherwise [5, 7]. It is also solvable in polynomial time on threshold
 125 graphs [5]. On bridgeless chordal graphs, there is a linear-time $(3/2)$ -approximation algorithm
 126 for RC, however the problem cannot be approximated with a factor less than $5/4$ on this
 127 graph class, unless $P = NP$ [6]. Some lower bounds on algorithms for solving RC are given
 128 by Kowalik et al. [17] and Agrawal [1] under the Exponential Time Hypothesis.

129 For the strong edge variant, an edge-colored graph is said to be *strongly rainbow-connected*
 130 if there is a rainbow shortest path between every pair of its vertices. The problem of deciding
 131 whether the edges of a given graph G can be colored in k colors to make G strongly rainbow-
 132 connected is referred to as SRC. For $k = 2$, it is not difficult to verify that RC is equivalent

133 to SRC. Not surprisingly, SRC is also NP-complete for $k \geq 2$ [2]. In contrast to RC, SRC
 134 remains hard on split graphs for every $k \geq 2$ [19, Theorem 4.1]. Moreover, on n -vertex split
 135 graphs, it is NP-hard to approximate SRC within a factor of $n^{1/2-\varepsilon}$ for any $\varepsilon > 0$, while
 136 RC admits an additive-1 approximation [5]. The former statement also holds for n -vertex
 137 bipartite graphs instead of split graphs [2]. For block graphs, computing SRC can be done in
 138 linear time [16], while RC on block graphs is conjectured to be hard (see [19, Conjecture 6.3]
 139 or [16]). In general, it appears that despite the interest, there are fewer complexity-theoretic
 140 results on SRC. In fact, the same is true when considering combinatorial results (see [25] for
 141 a broader discussion).

142 2 Preliminaries

143 In this paper, we work on undirected simple graphs. Such a graph is denoted by $G = (V, E)$,
 144 where V is the vertex set of G , and E is the edge set. We let n denote the number of vertices
 145 of G . For a vertex $x \in V$, $N(x)$ is the set of its *neighbors*, and $\deg(x) = |N(x)|$ is its *degree*.
 146 For a $S \subseteq V$, the subgraph of G induced by S is denoted by $G[S]$. A *cut vertex* of G is a
 147 vertex whose removal increases the number of connected components of G .

148 Given a path $P = x_1, x_2, \dots, x_{p-1}, x_p$ in G , the vertices from x_2 to x_{p-1} are called the
 149 *internal vertices* of P . The *distance* between two vertices u and v in G , denoted by $\text{dist}(u, v)$,
 150 is the length of a shortest path between u and v . The *diameter* of G , denoted by $\text{diam}(G)$,
 151 is the maximum distance between any pair of vertices of G .

152 A k -*coloring* of G is a function $c : V \rightarrow \{1, 2, \dots, k\}$. (From now on, we will denote a set
 153 of consecutive integers from 1 to k as $[k]$.) A *coloring* is simply a k -coloring for some $k \leq n$.
 154 A coloring c is *proper* if $c(u) \neq c(v)$ for every edge $uv \in E$. The *chromatic number* of G ,
 155 denoted by $\chi(G)$, is the smallest k such that G has a proper k -coloring. A d -*distance coloring*
 156 of G is a coloring c of G such that $c(u) \neq c(v)$ whenever $\text{dist}(u, v) \leq d$. The minimum
 157 number of colors needed for a d -distance coloring of G is known as the d -*distance chromatic*
 158 *number* of G , and it is denoted by $\chi_d(G)$. Note that $\chi_d(G)$ is equivalent to $\chi(G^d)$, i.e., the
 159 chromatic number of the d^{th} power of G .

160 Since, in this paper, we will only be working on the vertex variant of the rainbow coloring
 161 and rainbow connectivity, we might sometimes omit the word “vertex” when there is no
 162 confusion. The parameter $\text{srvc}(G)$ was defined by Li et al. [24], and they also verified that
 163 $\text{diam}(G) - 1 \leq \text{rvc}(G) \leq \text{srvc}(G) \leq n - 2$. The following upper bound was mentioned in [19]
 164 (see the same reference for further discussion and examples).

► **Proposition 5** ([19]). Let G be a connected graph with $\text{diam}(G) = d \geq 3$. Then

$$d - 1 \leq \text{rvc}(G) \leq \text{srvc}(G) \leq \chi_{d-2}(G).$$

165 **Proof.** There are at least two vertices in G connected by a shortest path of length d . Clearly,
 166 every coloring must use at least $d - 1$ colors to rainbow-connect this pair. On the other hand,
 167 between every pair of vertices u and v , there is a path of length at most d , meaning that
 168 it contains at most $d - 1$ internal vertices. As every $(d - 2)$ -distance coloring colors these
 169 internal vertices distinctly, the statement follows. ◀

170 A *dominating set* of G is a set $D \subseteq V$ such that every vertex in $V \setminus D$ is adjacent
 171 to at least one vertex in D . If $G[D]$ is connected, then D is a *connected dominating set*.
 172 The minimum size of a connected dominating set in G , denoted by $\gamma_c(G)$, is known as the
 173 *connected domination number* of G . This parameter provides an upper bound on the rainbow
 174 vertex connection number of a connected graph, since G becomes rainbow vertex-connected

175 by simply coloring all vertices of the connected dominating set distinctly, and the remaining
 176 vertices with any of the already used colors. This observation can be derived from [18].

177 ▶ Proposition 6 ([18]). If G is a connected graph, then $\mathbf{rvc}(G) \leq \gamma_c(G)$.

178 2.1 Graph classes

179 As we will be studying the mentioned problems on some graph classes, let us give a brief
 180 definition of these classes here. More definitions and properties will be added as needed
 181 when we handle these graphs. A detailed background on these graph classes can be found,
 182 for example, in the book by Brandstädt, Le, and Spinrad [3].

183 A graph is an *apex graph* if it contains a vertex (called an *apex*) whose removal results in
 184 a planar graph. A graph is *chordal* if all of its induced simple cycles are of length 3. Some
 185 well-known subclasses of chordal graphs are interval graphs, split graphs, and block graphs.
 186 A graph is an *interval graph* if it is chordal and it contains no triple of non-adjacent vertices,
 187 such that there is a path between every two of them that does not contain a neighbor of
 188 the third. A graph is a *split graph* if its vertex set can be partitioned into an independent
 189 set and a clique. A graph is a *block graph* if every biconnected component (block) of G is a
 190 complete graph.

191 Let σ be a permutation of the integers between 1 and n . We can make a graph G_σ
 192 on vertex set $[n]$ in the following way. Vertices i and j are adjacent in G_σ if and only if
 193 they appear in σ in the opposite order of their natural order. A graph on n vertices is a
 194 *permutation graph* if it is isomorphic to G_σ for some permutation σ of the integers between
 195 1 and n . A graph is a *bipartite permutation graph* if it is both a bipartite graph and a
 196 permutation graph.


197 2.2 Hypergraph coloring

198 For our hardness reductions we will use a well-known NP-complete problem called HYPER-
 199 GRAPH COLORING. A *hypergraph* $H = (N, \mathcal{E})$ with vertex set N and hyperedge set \mathcal{E} is a
 200 generalization of a graph, in which edges can contain more than two vertices. Thus \mathcal{E} consists
 201 of subsets of N of arbitrary size. The definition of a (vertex) coloring of a hypergraph is ex-
 202 actly that same as that of a graph. In a colored hypergraph, an edge is called *monochromatic*
 203 if all of its vertices received the same color. A *proper coloring* of a hypergraph generalizes a
 204 proper coloring of a graph in a natural way: we require that no hyperedge is monochromatic.
 205 To avoid trivial cases, we can assume from now on that every hyperedge contains at least
 206 two vertices. Thus a proper coloring must always use at least two colors.

207 The HYPERGRAPH COLORING problem takes as input a hypergraph H and an integer
 208 k and asks whether there is a proper coloring of H with at most k colors. The problem is
 209 well-known to be NP-complete for every $k \geq 2$ [26]. The GRAPH COLORING problem takes
 210 as input an undirected graph G and asks to determine the smallest k such that G has a
 211 proper k -coloring. This problem is NP-hard to approximate within a factor of $n^{1-\varepsilon}$ for any
 212 $\varepsilon > 0$, where n is the number of vertices [30]. Finally, the PLANAR 3-COLORING problem
 213 takes as input a planar graph G and asks whether G has a proper 3-coloring. This problem
 214 is NP-complete [14].

215 3 Bipartite graphs and their subclasses

216 In this section, we show that RVC and SRVC are hard on bipartite graphs for $k \geq 3$. We
 217 complement these results by showing that both problems can be solved in linear time on



■ **Figure 1** A hypergraph $H = (N, \mathcal{E})$ (left) transformed into a bipartite graph G (right) as described in the proof of Lemma 8. The dashed rectangle with rounded corners contains the sets in N' .

218 bipartite permutation graphs. We first observe that computing $\mathbf{rvc}(G)$ or $\mathbf{srvc}(G)$ is easy
 219 on bipartite graphs of diameter 3. The same observation was made by Li et al. [23].

220 ► **Proposition 7** ([23]). If G is a bipartite graph with $\text{diam}(G) = 3$, then $\mathbf{rvc}(G) = \mathbf{srvc}(G) = 2$.
 221 Moreover, such a coloring can be found in linear time.

222 **Proof.** The statement follows from Proposition 5 and the fact that every bipartite graph
 223 has a proper 2-coloring that can be found in linear time. ◀

224 It turns out that if $\text{diam}(G) \geq 4$, then $\mathbf{rvc}(G)$ and $\mathbf{srvc}(G)$ of a bipartite graph G
 225 become much harder to compute, as claimed in Theorem 2. We prove the following general
 226 construction.

227 ► **Lemma 8.** *Let H be a hypergraph on n vertices. Then in polynomial time we can construct
 228 a bipartite graph G of diameter 4 and with $O(n^3)$ vertices such that for any $k \in [n]$, H has a
 229 proper k -coloring if and only if G has a $(k + 1)$ -coloring under which G is (strongly) rainbow
 230 vertex-connected. Moreover, if H is a planar graph, then G is an apex graph.*

231 **Proof.** Let $H = (N, \mathcal{E})$ be an arbitrary hypergraph and let $n = |N|$. We construct a
 232 bipartite graph $G = (\{a\} \cup N' \cup I', E)$ where $N' = N'_1 \cup \dots \cup N'_{n+1}$, $I' = I'_1 \cup \dots \cup I'_{n+1}$,
 233 $N'_i := \{v_i \mid v \in N\}$, $I'_i := \{x_e^i \mid e \in \mathcal{E}\}$ and $E := \{av_i \mid v \in N, i \in [n+1]\} \cup \{v_i x_e^i \mid v \in N, e \in$
 234 $\mathcal{E}, i \in [n+1], v \in e\}$. Let $V = \{a\} \cup N' \cup I'$. A bipartition of G is given by $(\{a\} \cup I', N')$.
 235 Observe that $\text{diam}(G) = 4$ and that G has $O(n^3)$ vertices. Moreover, if H is a planar graph,
 236 then G consists of vertex a plus $n + 1$ copies of the graph obtained from H by subdividing
 237 each edge of H , and thus G is an apex graph. For an illustration of the construction, see
 238 Figure 1.

239 Consider any proper k -coloring $h : N \rightarrow [k]$ of H , i.e., no hyperedge of H is monochromatic
 240 under h . We construct a coloring $c : V \rightarrow [k + 1]$ in the following way. First, for every $v \in N$,
 241 we give the vertices v_1, v_2, \dots, v_n of G the same color as v , i.e., $c(v_i) = h(v)$ for all $v \in N$ and
 242 $i \in [n + 1]$. We give vertex a the color $k + 1$, i.e., $c(a) = k + 1$. The vertices in I all receive the
 243 same color, which is any arbitrary color in $[k + 1]$. Now we prove that G is strongly rainbow
 244 vertex-connected under c by showing that there is a rainbow vertex shortest path between

245 every pair of vertices. The only non-trivial case is when both vertices of the pair are in I .
 246 Consider two distinct vertices $x_e^i, x_f^j \in I$ (it is possible that $e = f$ or $i = j$ but not both).
 247 Since e and f are not monochromatic under h , we can pick two distinct vertices $u \in e$ and
 248 $v \in f$ such that $h(u) \neq h(v)$. It is clear that the path $x_e^i u a v x_f^j$ is a shortest path between x_e^i
 249 and x_f^j and that it is a rainbow vertex path. Hence, G is strongly rainbow vertex-connected
 250 under c .

251 Conversely, let c be a $(k + 1)$ -coloring of G under which G is (strongly) rainbow vertex-
 252 connected. For each $i \in [n + 1]$, define h_i to be the vertex coloring of H such that $h_i(v) = c(v_i)$
 253 for all $v \in N$. Let M_i be the set of vertices $v \in N$ such that $h_i(v) \neq c(a)$. Let $h'_i(v) = h_i(v)$
 254 if $v \in M_i$ and $h'_i(v) = 1$ otherwise. We claim that there exists an $i \in [n + 1]$ such that h'_i
 255 is a proper k -coloring of H . For the sake of contradiction, suppose that h'_i is not a proper
 256 k -coloring of H for every $i \in [n + 1]$. For each $i \in [n + 1]$, let $e_i \in \mathcal{E}$ be a monochromatic
 257 edge under h'_i . Suppose that, for some $i \in [n + 1]$, all vertices in e_i are colored $c(a)$ under
 258 c . Then any path from $x_{e_i}^i$ to $x_{e_i}^j$ for some $j \neq i$ uses two vertices having color $c(a)$ under
 259 c . Hence, c would not be a rainbow vertex coloring, a contradiction. Therefore, for each
 260 $i \in [n + 1]$, there is a vertex $v_i \in e_i$ for which $c(v_i) \neq c(a)$. Suppose now that for every
 261 $i \in [n + 1]$, all vertices in e_i are colored either $c(v_i)$ or $c(a)$ under c . If $c(v_i) = c(v_j)$ for
 262 $i \neq j$, then any path from $x_{e_i}^i$ to $x_{e_j}^j$ uses either two vertices having color $c(a)$ or two vertices
 263 having color $c(v_i) = c(v_j)$ under c . This would contradict the assumption that G is rainbow
 264 vertex-connected under c . Hence, $c(v_i) \neq c(v_j)$ for all distinct $i, j \in [n + 1]$. This implies that
 265 c uses at least $n + 2$ colors, a contradiction to the assumptions that c is a $(k + 1)$ -coloring
 266 of G and that $k \in [n]$. Therefore, for some $i \in [n + 1]$, there is a vertex $v'_i \in e_i$ for which
 267 $c(v'_i) \neq c(a)$ and $c(v'_i) \neq c(v_i)$. The latter implies that e_i is not monochromatic under h'_i , a
 268 contradiction. The claim follows, and thus H has a proper k -coloring. ◀

269 **Proof of Theorem 2.** For membership in NP, a certificate that $\text{rvc}(G) \leq k$ ($\text{srvc}(G) \leq k$)
 270 consists of a k -coloring and a list of (shortest) paths, one for every pair of non-adjacent vertices,
 271 that are rainbow vertex connected. For NP-hardness, we observe that the transformation
 272 of Lemma 8 implies a straightforward reduction from HYPERGRAPH COLORING. Since
 273 HYPERGRAPH COLORING is NP-complete for each $k \geq 2$, this proves the first part of the
 274 theorem.

275 For the second part of the theorem, we consider an instance of GRAPH COLORING that
 276 consists of a graph on ℓ vertices and apply Lemma 8. Note that the total number of vertices
 277 in G is $n = O(\ell^3)$. From the hardness of approximation of GRAPH COLORING, we know that
 278 for all $\varepsilon > 0$, it is NP-hard to distinguish between the case when H is properly colorable with
 279 ℓ^ε colors and the case when H is not properly colorable with fewer than $\ell^{1-\varepsilon}$ colors [30]. By
 280 Lemma 8, this implies that it is NP-hard to distinguish between the case when G is (strong)
 281 rainbow vertex colorable with $\ell^\varepsilon + 1 \leq n^\varepsilon + 1$ colors and the case when G is not (strong)
 282 rainbow vertex colorable with fewer than $\ell^{1-\varepsilon} + 1 = \Omega(n^{1/3-\varepsilon})$ colors. The second statement
 283 of the theorem follows. ◀

284 We then proceed to give a proof of Theorem 1. This result can be considered as a first
 285 step to understand rainbow coloring on sparse graphs classes.

286 **Proof of Theorem 1.** The proof follows along the same lines as the proof of the first part
 287 of Theorem 2. Instead of HYPERGRAPH COLORING, however, we reduce from PLANAR
 288 3-COLORING, the problem of deciding whether a planar graph has a proper 3-coloring. This
 289 problem is NP-complete. The statement follows from Lemma 8, because the graph resulting
 290 from the construction is a bipartite apex graph of diameter 4.

291 For the hardness of approximation, we recall that any planar graph has a proper 4-coloring,
 292 and thus the graph G constructed in Lemma 8 has a 5-coloring under which G is rainbow
 293 vertex-connected. Hence, Lemma 8 combined with the NP-hardness of PLANAR 3-COLORING
 294 makes it NP-hard to decide whether G has a 5-coloring or a 4-coloring under which G is
 295 rainbow vertex-connected. ◀

296 We now complement the above hardness results with a positive result in the case when a
 297 bipartite graph is also a permutation graph, as claimed in Theorem 4. Bipartite permutation
 298 graphs have a desirable property, related to breadth-first search (BFS), that we will use
 299 heavily in our next result. Let us first define a chain graph. A bipartite graph is a *chain graph*
 300 if the vertices of the two independent sets A and B can be ordered as $\{a_1, a_2, \dots, a_k\}$ and
 301 $\{b_1, b_2, \dots, b_\ell\}$, such that $N(a_1) \subseteq N(a_2) \subseteq \dots \subseteq N(a_k)$, equivalently, $N(b_\ell) \subseteq N(b_{\ell-1}) \subseteq$
 302 $\dots \subseteq N(b_1)$.

303 In every bipartite permutation graph G it is possible to find a vertex v such that the levels
 304 L_0, L_1, L_2, \dots of the tree resulting from a BFS starting from v have the following properties.
 305 For all i , $L_0 = \{v\}$, L_i is an independent set and $G[L_i \cup L_{i+1}]$ is a chain graph. Moreover, for
 306 each level i , there exists a special vertex $a_i \in L_i$ such that $L_{i+1} \subset N(a_i)$. The vertex v can
 307 be picked as the first vertex of a *strong ordering*. It has been shown by Spinrad et al. [27]
 308 that a bipartite graph is a permutation graph if and only if it has a strong ordering, and
 309 such an ordering can be computed in linear time. The properties of the BFS tree above are
 310 well-known and easy to deduce from a strong ordering [29].

311 ► **Theorem 9.** *If G is a bipartite permutation graph, then $\text{rvc}(G) = \text{srvc}(G) = \text{diam}(G) - 1$,
 312 and the corresponding (strong) rainbow vertex coloring can be found in time that is linear in
 313 the size of G .*

314 **Proof.** Let $G = (V, E)$ be a bipartite permutation graph. Let v be a first vertex in a strong
 315 ordering for G . We start by doing a BFS on G with v as the root. Let k be the number
 316 of levels in the BFS tree in addition to level 0. Hence, L_i is the set of vertices in level
 317 i of the BFS tree, $0 \leq i \leq k$, with $L_0 = \{v\}$. Since $\text{dist}(v, y) = k$ for every $y \in L_k$, we
 318 conclude that $\text{diam}(G) \geq k$. Furthermore, if $\text{dist}(x, y) > k - 1$ for some $x \in L_1$ and some
 319 $y \in L_k$, then we can conclude that $\text{dist}(x, y) = k + 1$, where $x, v, a_1, a_2, \dots, a_{k-1}, y$ is a shortest
 320 path between x and y . In this case, $\text{diam}(G) = k + 1$. We distinguish between these two cases:

321

322 **Case 1.** $\text{diam}(G) = k$.

323 We construct a strong rainbow vertex coloring $c : V \rightarrow [k - 1]$ for G in the following way.
 324 If $x \in L_i$, we define $c(x) = i$, for $1 \leq i \leq k - 1$. We define $c(v) = k - 1$, and we give arbitrary
 325 colors between 1 and $k - 1$ to the vertices of L_k . To see that G is indeed rainbow-connected
 326 under c , consider any pair $x, y \in V$. If $xy \in E$ or if they are in the same level of the BFS
 327 tree, there is nothing to prove, since $\text{dist}(x, y) \leq 2$. Otherwise, we have exactly the following
 328 cases:

- 329 1. $x = v$ and $y \in L_j$: Then the path $v, a_1, \dots, a_{j-1}, y$ is shortest and it is rainbow.
- 330 2. $x \in L_1$ and $y \in L_k$: In this case, $\text{dist}(x, y) = k - 1$. Otherwise, since each L_i is an
 331 independent set, we would have $\text{dist}(x, y) \geq k + 1$, which contradicts our assumption that
 332 $\text{diam}(G) = k$. Since $\text{dist}(x, y) = k - 1$, every shortest path between x and y is rainbow,
 333 as every vertex of such a shortest path has to be in a distinct level of the BFS tree.
- 334 3. $x \in L_1$ and $y \in L_j$ with $2 \leq j \leq k - 1$: If $\text{dist}(x, y) = j - 1$, then again by the same
 335 argument used above, every shortest path between x and y is rainbow. If $\text{dist}(x, y) > j - 1$,
 336 then $\text{dist}(x, y) = j + 1$, and the shortest path $x, v, a_1, \dots, a_{j-1}, y$ has distinct colors on

337 all its internal vertices. (Note that y might have the same color as v if $j = k - 1$, but this
338 is fine since y is the end of the path.)

339 **4.** $x \in L_i$ and $y \in L_j$ with $2 \leq i < j \leq k$: If $\text{dist}(x, y) = j - i$, then every shortest path
340 is rainbow. If $\text{dist}(x, y) > j - i$, then the path $x, a_{i-1}, a_i, \dots, a_{j-1}, y$ is rainbow and has
341 length $j - i + 1$, and it is therefore shortest.

342 **Case 2.** $\text{diam}(G) = k + 1$.

343 We construct a strong rainbow vertex coloring $c : V \rightarrow [k]$ for G in the following way. If
344 $x \in L_i$, we define $c(x) = i$, for $1 \leq i \leq k - 1$. We define $c(v) = k$, and we give arbitrary
345 colors between 1 and k to the vertices of L_k . To see that G is indeed rainbow-connected
346 under c , consider any pair $x, y \in V$. Again, if $xy \in E$ or if they are in the same level of the
347 BFS,

348 case:

349 ■ $x \in L_i$ and $y \in L_j$, with $0 \leq i < j \leq k$: If $\text{dist}(x, y) = j - i$ then every shortest path
350 between x and y is rainbow. Otherwise, the path $x, a_{i-1}, a_i, \dots, a_{j-1}, y$ is rainbow and
351 has length $j - i + 2$, therefore being shortest.

352 In both cases, c is a strong rainbow vertex coloring for G with $\text{diam}(G) - 1$ colors. By
353 Proposition 5 we can conclude that $\text{rvc}(G) = \text{srvc}(G) = \text{diam}(G) - 1$. ◀

354 **4 Chordal graphs and their subclasses**

355 In this section, we investigate the complexity of RVC and SRVC on chordal graphs and
356 some subclasses of chordal graphs. We start by proving that both problems are NP-complete
357 when the input graph is a split graph, implying that they are also NP-complete on chordal
358 graphs. On the positive side, we show that RVC is polynomial-time solvable on interval
359 graphs, and both RVC and SRVC are polynomial-time solvable on block graphs and on unit
360 interval graphs.

361 We start by observing that computing $\text{rvc}(G)$ or $\text{srvc}(G)$ is easy on graphs of diameter 2.

362 ▶ **Proposition 10** ([18]). If G is a graph with $\text{diam}(G) = 2$, then $\text{rvc}(G) = \text{srvc}(G) = 1$.
363 Moreover, such a coloring can be found in linear time.

364 **Proof.** Color each vertex of G with the same color. Since each shortest path between two
365 vertices contains at most one internal vertex, G is strongly rainbow vertex-connected under
366 this coloring. ◀

367 If G is a split graph of $\text{diam}(G) = 3$ (note that split graphs have diameter at most 3),
368 then $\text{rvc}(G)$ and $\text{srvc}(G)$ become much harder to compute, as claimed in Theorem 3. We
369 prove the following general construction, which closely mimics the construction of Lemma 8.

370 ▶ **Lemma 11.** *Let H be a hypergraph on n vertices. Then in polynomial time we can
371 construct a split graph G of diameter 3 and with $O(n^3)$ vertices such that for any $k \in [n]$, H
372 has a proper k -coloring if and only if G has a k -coloring under which G is (strongly) rainbow
373 vertex-connected.*

374 **Proof.** Let $H = (N, \mathcal{E})$ be an arbitrary hypergraph and let $n = |N|$. We construct a split
375 graph $G = (N' \cup I', E)$ where $N' = N'_1 \cup \dots \cup N'_{n+1}$, $I' = I'_1 \cup \dots \cup I'_{n+1}$, $N'_i := \{v_i \mid v \in N\}$,
376 $I'_i := \{x_e^i \mid e \in \mathcal{E}\}$ and $E := \{u_i v_j \mid u, v \in N, i, j \in [n + 1]\} \cup \{v_i x_e^i \mid v \in N, e \in \mathcal{E}, i \in$
377 $[n + 1], v \in e\}$. Let $V = N' \cup I'$. The constructed graph G is a split graph since $G[I']$ is an
378 independent set and $G[N']$ is a clique. Observe that $\text{diam}(G) = 3$ and that G has $O(n^3)$

379 vertices. The construction is illustrated in Figure 1: note that since $G[N']$ is a clique, all
 380 possible edges now appear between the vertices inside the rectangle with rounded corners.

381 Consider any proper k -coloring $h : N \rightarrow [k]$ of H , i.e., no hyperedge of H is monochromatic
 382 under h . We construct a coloring $c : V \rightarrow [k]$ in the following way. First, for every $v \in N$,
 383 we give the vertices v_1, v_2, \dots, v_n of G the same color as v , i.e., $c(v_i) = h(v)$ for all $v \in N$
 384 and $i \in [n + 1]$. The vertices in I all receive the same color, which is any arbitrary color in
 385 $[k]$. Now, we prove that G is strongly rainbow vertex-connected under c by showing that
 386 there is a rainbow vertex shortest path between every pair of vertices. The only non-trivial
 387 case is when both vertices of the pair are in I . Consider two distinct vertices $x_e^i, x_f^j \in I$ (it is
 388 possible that $e = f$ or $i = j$ but not both). Since e and f are not monochromatic under h ,
 389 we can pick two distinct vertices $u \in e$ and $v \in f$ such that $h(u) \neq h(v)$. It is clear that the
 390 path $x_e^i u v x_f^j$ is a shortest path between x_e^i and x_f^j and that it is rainbow vertex path.

391 Conversely, let c be a k -coloring of G under which G is (strongly) rainbow vertex-connected.
 392 For each $i \in [n + 1]$, define h_i to be the vertex coloring of H such that $h_i(v) = c(v_i)$ for
 393 all $v \in N$. We claim that there exists an $i \in [n + 1]$ such that h_i is a proper k -coloring of
 394 H . For the sake of contradiction, suppose that h_i is not a proper k -coloring of H for every
 395 $i \in [n + 1]$. For each $i \in [n + 1]$, let $e_i \in \mathcal{E}$ be a monochromatic edge under h_i . Let v_i be an
 396 arbitrary vertex in e_i . Suppose now that for every $i \in [n + 1]$, all vertices in e_i are colored
 397 $c(v_i)$ under c . If $c(v_i) = c(v_j)$ for $i \neq j$, then any path from $x_{e_i}^i$ to $x_{e_j}^j$ uses two vertices
 398 having color $c(v_i) = c(v_j)$ under c . This would contradict the assumption that G is rainbow
 399 vertex-connected under c . Hence, $c(v_i) \neq c(v_j)$ for all distinct $i, j \in [n + 1]$. This implies
 400 that c uses at least $n + 1$ colors, a contradiction to the assumptions that c is a k -coloring of
 401 G and $k \in [n]$. Therefore, for some $i \in [n + 1]$, there is a vertex $v'_i \in e_i$ for which $c(v'_i) \neq c(a)$
 402 and $c(v'_i) \neq c(v_i)$. The latter implies that e_i is not monochromatic under h'_i , a contradiction.
 403 The claim follows, and thus H has a proper H -coloring. ◀

404 **Proof of Theorem 3.** The proof follows in exactly the same way as Theorem 2, except that
 405 we apply Lemma 11 instead of Lemma 8. ◀

406 We now move on to the positive results. As a consequence of the following theorems, we
 407 complete the proof of Theorem 4.

408 **► Theorem 12.** *Let G be a block graph, and let ℓ be the number of cut vertices in G . Then*
 409 $\mathbf{rvc}(G) = \mathbf{srvc}(G) = \ell$. *The corresponding (strong) rainbow vertex coloring can be found in*
 410 *time that is linear in the size of G .*

411 **Proof.** Let $G = (V, E)$ be a block graph and $\{a_1, a_2, \dots, a_\ell\}$ be the set of cut vertices of
 412 G . We construct a strong rainbow vertex coloring $c : V \rightarrow [\ell]$ for G by defining $c(a_i) = i$
 413 for $i \in [\ell]$ and giving the other vertices arbitrary colors between 1 and ℓ . An important
 414 property of block graphs is that there is a unique shortest path between every pair of vertices.
 415 Moreover, each internal vertex of such a path is a cut vertex. Since all the cut vertices
 416 received distinct colors, these shortest paths are all rainbow. The proof follows by observing
 417 that $\mathbf{rvc}(G) \geq \mathbf{srvc}(G) \geq \ell$ as well. ◀

418 For our next result, we need to mention that every interval graph has a representation
 419 called an *interval model*. Let \mathcal{I} be a set of n intervals of the real line. Then we can define
 420 a graph $G_{\mathcal{I}}$ with a vertex for each interval, such that two vertices are adjacent if and only
 421 if their corresponding intervals overlap. A graph G is an interval graph if and only if G is
 422 isomorphic to $G_{\mathcal{I}}$ for some set \mathcal{I} of intervals. In this case \mathcal{I} is called an interval model of G .

423 ► **Theorem 13.** *If G is an interval graph, then $\text{rvc}(G) = \text{diam}(G) - 1$, and the corresponding*
 424 *rainbow vertex coloring can be found in time that is linear in the size of G .*

425 **Proof.** Let $G = (V, E)$ be an interval graph and \mathcal{I} be an interval model for G . The interval
 426 corresponding to vertex v is denoted by I_v . For each interval $I \in \mathcal{I}$, we let $r(I)$ be its right
 427 endpoint and $\ell(I)$ its left endpoint. Let $I_u \in \mathcal{I}$ be such that $r(I_u) \leq r(I)$ for all $I \in \mathcal{I}$.
 428 Let $I_v \in \mathcal{I}$ be such that $\ell(I_v) \geq \ell(I)$ for all $I \in \mathcal{I}$. Let $P = u, x_1, x_2, \dots, x_k, v$ be a shortest
 429 path between u and v in G . Observe that P is a connected dominating set. Furthermore,
 430 since P is a shortest path, $k \leq \text{diam}(G) - 1$. By the way we defined u and v , we have
 431 that $N(u) \subseteq N(x_1)$ and $N(v) \subseteq N(x_k)$. This implies that the set $\{x_1, x_2, \dots, x_k\}$ is also a
 432 connected dominating set. By Proposition 6, G has a rainbow vertex coloring $c : V \rightarrow [k]$
 433 with $c(x_i) = i$, and we can give all the other vertices arbitrary colors. ◀

434 An interval graph is a *unit interval graph* if it has an interval model in which every interval
 435 has the same length (or no interval properly contains another interval). Unit interval graphs
 436 have the same BFS tree structure as that of bipartite permutation graphs, with the single
 437 difference that every level of the BFS tree is a clique instead of an independent set [15].

438 ► **Theorem 14.** *If G is a unit interval graph, then $\text{rvc}(G) = \text{srvc}(G) = \text{diam}(G) - 1$, and*
 439 *the corresponding (strong) rainbow vertex coloring can be found in time that is linear in the*
 440 *size of G .*

441 **Proof.** Let $G = (V, E)$ be a unit interval graph. Let v be the vertex corresponding to a
 442 first interval in an ordering of the intervals in the unit interval model of G by their right
 443 endpoints. Do a BFS on G with v as the root. Let L_i be the set of vertices in level i of the
 444 BFS tree, $0 \leq i \leq k$, with $L_0 = \{v\}$. Recall that, for $0 \leq i \leq k - 1$, there exists a special
 445 vertex $a_i \in L_i$ such that $L_{i+1} \subset N(a_i)$.

446 Consider a vertex $u \in L_k$. A shortest path between v and u has $k - 1$ internal vertices,
 447 which implies that $\text{diam}(G) \geq k$. To construct a strong rainbow coloring $c : V \rightarrow [k - 1]$, we
 448 assign, for $1 \leq i \leq k - 1$, $c(x) = i$ if $x \in L_i$ and we give arbitrary colors to the vertices of L_k .

449 To see that G is strongly rainbow vertex-connected under c , consider $x, y \in V$. If both x
 450 and y are in the same level of the BFS tree, then they are adjacent. So let us consider the
 451 case when $x \in L_i$ and $y \in L_j$, with $1 \leq i < j \leq k$. If there is a shortest path between x and
 452 y each of whose vertices is in a distinct level of the BFS tree, then this path is rainbow. If
 453 this is not the case, we consider the path $x, a_i, a_{i+1}, \dots, a_{j-1}, y$. In this case, this path is a
 454 shortest path between x and y , and its internal vertices have distinct colors, since only x
 455 and a_i belong to the same level of the BFS. This proves that c is indeed a strong rainbow
 456 coloring for G with $\text{diam}(G) - 1$ colors. ◀

457 5 Concluding remarks and related problems

458 It should be mentioned that other variants of rainbow problems have been studied as well.
 459 When a coloring of the edges or the vertices of a graph is already given as input, we can
 460 ask whether the graph is rainbow-connected or rainbow vertex-connected. Both of these
 461 problems are known to be NP-complete even on highly restricted graphs, like interval graphs,
 462 series-parallel graphs, and k -regular graphs for every $k \geq 3$ [21, 20, 28]. However, we stress
 463 that these problems are strictly different from RC and RVC. That is, complexity results on
 464 one problem are not transferable to the other.

465 Finally, we end our paper with the following open question.¹ A *diametral path* of a graph
 466 G is a shortest path whose length is equal to $\text{diam}(G)$. A graph is a *diametral path* if every
 467 connected induced subgraph has a dominating diametral path.

468 ► **Conjecture 15.** Let G be a diametral path graph. Then $\text{rvc}(G) = \text{diam}(G) - 1$.

469 — References —

- 470 **1** Akanksha Agrawal. Fine-grained complexity of rainbow coloring and its variants. In *Pro-*
 471 *ceedings of the 42nd International Symposium on Mathematical Foundations of Computer*
 472 *Science (MFCS)*, pages 60:1–60:14, 2017.
- 473 **2** Prabhanjan Ananth, Meghana Nasre, and Kanthi K. Sarpatwar. Rainbow connectivity:
 474 Hardness and tractability. In *IARCS Annual Conference on Foundations of Software Tech-*
 475 *nology and Theoretical Computer Science (FSTTCS)*, pages 241–251, 2011.
- 476 **3** Andreas Brandstädt, Van Bang Le, and Jeremy P. Spinrad. *Graph Classes: A Survey*.
 477 SIAM Monographs in Discrete Mathematics and Applications, 1999.
- 478 **4** Sourav Chakraborty, Eldar Fischer, Arie Matsliah, and Raphael Yuster. Hardness and
 479 algorithms for rainbow connection. *Journal of Combinatorial Optimization*, 21(3):330–347,
 480 2011.
- 481 **5** L. Sunil Chandran and Deepak Rajendraprasad. Rainbow Colouring of Split and Threshold
 482 Graphs. In *Proceedings of the 18th Annual International Computing and Combinatorics*
 483 *Conference (COCOON)*, volume 7434 of *Lecture Notes in Computer Science*, pages 181–192.
 484 Springer, 2012.
- 485 **6** L. Sunil Chandran and Deepak Rajendraprasad. Inapproximability of rainbow colouring.
 486 In *IARCS Annual Conference on Foundations of Software Technology and Theoretical Com-*
 487 *puter Science (FSTTCS)*, pages 153–162, 2013.
- 488 **7** L. Sunil Chandran, Deepak Rajendraprasad, and Marek Tesař. Rainbow colouring of split
 489 graphs. *Discrete Applied Mathematics*, 216:98–113, 2017.
- 490 **8** Gary Chartrand, Garry L Johns, Kathleen A McKeon, and Ping Zhang. Rainbow connec-
 491 tion in graphs. *Mathematica Bohemica*, 133(1):85–98, 2008.
- 492 **9** Lily Chen, Xueliang Li, and Huishu Lian. Further hardness results on the rainbow vertex-
 493 connection number of graphs. *Theoretical Computer Science*, 481:18–23, 2013.
- 494 **10** Lily Chen, Xueliang Li, and Yongtang Shi. The complexity of determining the rainbow
 495 vertex-connection of a graph. *Theoretical Computer Science*, 412(35):4531–4535, 2011.
- 496 **11** Paul Dorbec, Ingo Schiermeyer, Elżbieta Sidorowicz, and Éric Sopena. Rainbow connection
 497 in oriented graphs. *Discrete Applied Mathematics*, 179:69–78, 2014.
- 498 **12** Eduard Eiben, Robert Ganian, and Juho Lauri. On the complexity of rainbow coloring
 499 problems. In *Proceedings of the 26th International Workshop on Combinatorial Algorithms*
 500 *(IWOCA)*, volume 9538 of *Lecture Notes in Computer Science*, pages 209–220. Springer,
 501 2015.
- 502 **13** Eduard Eiben, Robert Ganian, and Juho Lauri. On the complexity of rainbow coloring
 503 problems. *Discrete Applied Mathematics*, 246:38–48, 2018.
- 504 **14** M. R. Garey, David S. Johnson, and Larry J. Stockmeyer. Some Simplified NP-Complete
 505 Graph Problems. *Theoretical Computer Science*, 1(3):237–267, 1976.
- 506 **15** Pavol Hell and Jing Huang. Certifying LexBFS Recognition Algorithms for Proper Interval
 507 Graphs and Proper Interval Bigraphs. *SIAM Journal on Discrete Mathematics*, 18(3):554–
 508 570, 2004.

¹ The statement is claimed in [19, Proposition 5.2] but its proof contains an error. Essentially, only an upper bound of $\text{diam}(G) + 1$ is known by Proposition 6.

- 509 **16** Melissa Keranen and Juho Lauri. Computing minimum rainbow and strong rainbow color-
510 ings of block graphs. *arXiv preprint arXiv:1405.6893*, 2014.
- 511 **17** Łukasz Kowalik, Juho Lauri, and Arkadiusz Socała. On the fine-grained complexity of
512 rainbow coloring. In *Proceedings of the 24th Annual European Symposium on Algorithms*
513 (*ESA*), pages 58:1–58:16, 2016.
- 514 **18** Michael Krivelevich and Raphael Yuster. The rainbow connection of a graph is (at most)
515 reciprocal to its minimum degree. *Journal of Graph Theory*, 63(3):185–191, 2010.
- 516 **19** Juho Lauri. *Chasing the Rainbow Connection: Hardness, Algorithms, and Bounds*. PhD
517 thesis, Tampere University of Technology, 2016.
- 518 **20** Juho Lauri. Further hardness results on rainbow and strong rainbow connectivity. *Discrete*
519 *Applied Mathematics*, 201:191–200, 2016.
- 520 **21** Juho Lauri. Complexity of rainbow vertex connectivity problems for restricted graph classes.
521 *Discrete Applied Mathematics*, 219:132–146, 2017.
- 522 **22** Van Bang Le and Zsolt Tuza. Finding optimal rainbow connection is hard. Technical
523 Report CS-03-09, Universität Rostock, 2009.
- 524 **23** Shasha Li, Xueliang Li, and Yongtang Shi. Note on the complexity of deciding the rain-
525 bow (vertex-)connectedness for bipartite graphs. *Applied Mathematics and Computation*,
526 258:155–161, 2015.
- 527 **24** Xueliang Li, Yaping Mao, and Yongtang Shi. The strong rainbow vertex-connection of
528 graphs. *Utilitas Mathematica*, 93:213–223, 2014.
- 529 **25** Xueliang Li, Yongtang Shi, and Yuefang Sun. Rainbow connections of graphs: A survey.
530 *Graphs and Combinatorics*, 29(1):1–38, 2013.
- 531 **26** László Lovász. Coverings and colorings of hypergraphs. In *Proceedings of the 4th South-*
532 *eastern Conference on Combinatorics, Graph Theory, and Computing*, pages 3–12, 1973.
- 533 **27** Jeremy Spinrad, Andreas Brandstädt, and Lorna Stewart. Bipartite permutation graphs.
534 *Discrete Applied Mathematics*, 18(3):279–292, 1987.
- 535 **28** Kei Uchizawa, Takanori Aoki, Takehiro Ito, Akira Suzuki, and Xiao Zhou. On the Rainbow
536 Connectivity of Graphs: Complexity and FPT Algorithms. *Algorithmica*, 67(2):161–179,
537 2013.
- 538 **29** Ryuhei Uehara and Gabriel Valiente. Linear structure of bipartite permutation graphs and
539 the longest path problem. *Information Processing Letters*, 103(2):71–77, 2007.
- 540 **30** David Zuckerman. Linear degree extractors and the inapproximability of max clique and
541 chromatic number. In *Proceedings of the 38th Annual ACM Symposium on Theory of*
542 *Computing (STOC)*, pages 681–690. ACM, 2006.