

The Impact of Geometry on Monochrome Regions in the Flip Schelling Process

Thomas Bläsius  

Karlsruhe Institute of Technology, Germany

Tobias Friedrich   

Hasso Plattner Institute, University of Potsdam, Germany

Martin S. Krejca   

Sorbonne University, CNRS, LIP6, Paris, France

Louise Molitor   

Hasso Plattner Institute, University of Potsdam, Germany

Abstract

Schelling’s classical segregation model gives a coherent explanation for the wide-spread phenomenon of residential segregation. We introduce an agent-based saturated open-city variant, the Flip Schelling Process (FSP), in which agents, placed on a graph, have one out of two types and, based on the predominant type in their neighborhood, decide whether to change their types; similar to a new agent arriving as soon as another agent leaves the vertex.

We investigate the probability that an edge $\{u, v\}$ is monochrome, i.e., that both vertices u and v have the same type in the FSP, and we provide a general framework for analyzing the influence of the underlying graph topology on residential segregation. In particular, for two adjacent vertices, we show that a highly decisive common neighborhood, i.e., a common neighborhood where the absolute value of the difference between the number of vertices with different types is high, supports segregation and, moreover, that large common neighborhoods are more decisive.

As an application, we study the expected behavior of the FSP on two common random graph models with and without geometry: (1) For random geometric graphs, we show that the existence of an edge $\{u, v\}$ makes a highly decisive common neighborhood for u and v more likely. Based on this, we prove the existence of a constant $c > 0$ such that the expected fraction of monochrome edges after the FSP is at least $1/2 + c$. (2) For Erdős–Rényi graphs we show that large common neighborhoods are unlikely and that the expected fraction of monochrome edges after the FSP is at most $1/2 + o(1)$. Our results indicate that the cluster structure of the underlying graph has a significant impact on the obtained segregation strength.

2012 ACM Subject Classification Theory of computation → Network formation; Theory of computation → Random network models

Keywords and phrases Agent-based Model, Schelling Segregation, Spin System

Digital Object Identifier 10.4230/LIPIcs.ISAAC.2021.29

Related Version *Full Version:* <https://arxiv.org/pdf/2102.09856.pdf> [12]

Funding *Martin S. Krejca:* This work was supported by the Paris Île-de-France Region.

Acknowledgements We want to thank Thomas Sauerwald for the discussions on random walks.

1 Introduction

Residential segregation is a well-known sociological phenomenon [49] where different groups of people tend to separate into largely homogeneous neighborhoods. Studies, e.g., [18], show that individual preferences are the driving force behind present residential patterns and bear much to the explanatory weight. Local choices therefore lead to a global phenomenon [47]. A simple model for analyzing residential segregation was introduced by Schelling [46, 47] in



© Thomas Bläsius, Tobias Friedrich, Martin S. Krejca, and Louise Molitor;
licensed under Creative Commons License CC-BY 4.0

32nd International Symposium on Algorithms and Computation (ISAAC 2021).

Editors: Hee-Kap Ahn and Kunihiko Sadakane; Article No. 29; pp. 29:1–29:17

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

the 1970s. In his model, two types of agents, placed on a grid, act according to the following threshold behavior, with $\tau \in (0, 1)$ as the *intolerance threshold*: agents are *content* with their current position on the grid if at least a τ -fraction of neighbors is of their own type. Otherwise they are *discontent* and want to move, either via swapping with another random discontent agent or via jumping to a vacant position. Schelling demonstrated via simulations that, starting from a uniform random distribution, the described process drifts towards strong segregation, even if agents are tolerant and agree to live in mixed neighborhoods, i.e., if $\tau \leq \frac{1}{2}$. Many empirical studies have been conducted to investigate the influence of various parameters on the obtained segregation, see [8, 9, 25, 41, 45]. On the theoretical side, Schelling's model started recently gaining traction within the algorithmic game theory and artificial intelligence communities [1, 11, 16, 17, 21, 22, 33], with focus on core game theoretic questions, where agents strategically select locations. Henry et al. [31] described a simple model of graph clustering motivated by Schelling where they showed that segregated graphs always emerge. Variants of the random Schelling segregation process were analyzed by a line of work that showed that residential segregation occurs with high probability [5, 7, 10, 13, 32, 51].


We initiate the study of an agent-based model, called the *Flip Schelling Process (FSP)*, which can be understood as the Schelling model in a *saturated open city*. In contrast to *closed cities* [7, 13, 32, 51], which require fixed populations, open cities [4, 5, 10, 27] allow resident to move away. In saturated city models, also known as voter models [20, 35, 36], vertices are not allowed to be unoccupied, hence, a new agent enters as soon as one agent vacates a vertex. In general, in voter models, two types of agents are placed on a graph. Agents examine their neighbors and, if a certain threshold is of another type, they change their types. Also in this model, segregation is visible. There is a line of work, mainly in physics, that studies the voting dynamics on several types of graphs [3, 14, 37, 43, 50]. Related to voter models, Granovetter [30] proposed another threshold model treating binary decisions and spurred a number of research, which studied and motivated variants of the model, see [2, 34, 38, 44].

In the FSP, agents have binary types. An agent is content if the fraction of agents in its neighborhood with the same type is larger than $\frac{1}{2}$. Otherwise, if the fraction is smaller than $\frac{1}{2}$, an agent is discontent and is willing to flip its type to become content. If the fraction of same type agents in its neighborhood is exactly $\frac{1}{2}$, an agent flips its type with probability $\frac{1}{2}$. Starting from an initial configuration where the type of each agent is chosen uniformly at random, we investigate a simultaneous-move, one-shot process and bound the number of monochrome edges, which is a popular measurement for segregation strength [19, 26].

Close to our model is the work by Omidvar and Franceschetti [39, 40], who initiated an analysis of the size of monochrome regions in the so called *Schelling Spin Systems*. Agents of two different types are placed on a grid [39] and a geometric graph [40], respectively. Then independent and identical Poisson clocks are assigned to all agents and, every time a clock rings, the state of the corresponding agent is flipped if and only if the agent is discontent w.r.t. a certain intolerance threshold τ regarding the neighborhood size. The model corresponds to the Ising model with zero temperature with Glauber dynamics [15, 48].

The commonly used underlying topology for modeling the residential areas are (toroidal) grid graphs [11, 32, 39], regular graphs [11, 17, 21], paths [11, 33], cycles [4, 6, 7, 13, 51] and trees [1, 11, 22, 33]. Considering the influence of the given topology that models the residential area regarding, e.g., the existence of stable states and convergence behavior leads to phenomena like non-existence of stable states [21, 22], non-convergence to stable states [11, 17, 21], and high-mixing times in corresponding Markov chains [10, 28].

To avoid such undesirable characteristics, we suggest to investigate *random geometric graphs* [42], like in [40]. Random geometric graphs demonstrate, in contrast to other random graphs without geometry, such as *Erdős-Rényi graphs* [23, 29], community structures, i.e.,



■ **Figure 1** The fraction of monochrome edges after the Flip Schelling Process (FSP) in Erdős–Rényi graphs and random geometric graphs for different graph sizes (number of vertices n) and different expected average degrees. Each data point shows the average over 1000 generated graphs with one simulation of the FSP per graph. The error bars show the interquartile range, i.e., 50% of the measurements lie between the top and bottom end of the error bar.

densely connected clusters of vertices. An effect observed by simulating the FSP is that the fraction of monochrome edges is significantly higher in random geometric graphs compared to Erdős–Rényi graphs, where the fraction stays almost stable around $\frac{1}{2}$, cf. Fig 1.

We set out for rigorously proving this phenomenon. In particular, we prove for random geometric graphs that there exists a constant c such that, given an edge $\{u, v\}$, the probability that $\{u, v\}$ is monochrome is lower-bounded by $\frac{1}{2} + c$, cf. Theorem 6. In contrast, we show for Erdős–Rényi graphs that segregation is not likely to occur and that the probability that $\{u, v\}$ is monochrome is upper-bounded by $\frac{1}{2} + o(1)$, cf. Theorem 17.

We introduce a general framework to deepen the understanding of the influence of the underlying topology on residential segregation. To this end, we first show that a highly decisive common neighborhood supports segregation, cf. Section 3.1. In particular, we provide a lower bound on the probability that an edge $\{u, v\}$ is monochrome based on the probability that the difference between the majority and the minority regarding both types in the common neighborhood, i.e., the number of agents which are adjacent to u and v , is larger than their exclusive neighborhoods, i.e., the number of agents which are adjacent to either u or v . Next, we show that large sets of agents are more decisive, cf. Section 3.2. This implies that a large common neighborhood, compared to the exclusive neighborhood, is likely to be more decisive, i.e., makes it more likely that the absolute value of the difference between the number of different types in the common neighborhood is larger than in the exclusive ones. These considerations hold for arbitrary graphs. Hence, we reduce the question concerning a lower bound for the fraction of monochrome edges in the FSP to the probability that, given $\{u, v\}$, the common neighborhood is larger than the exclusive neighborhoods of u and v , respectively.

For random geometric graphs, we prove that a large geometric region, i.e., the intersecting region that is formed by intersecting disks, leads to a large vertex set, cf. Section 3.3, and that random geometric graphs have enough edges that have sufficiently large intersecting regions, cf. Section 3.4, such that segregation is likely to occur. In contrast, for Erdős–Rényi graphs, we show that the common neighborhood between two vertices u and v is with high probability empty and therefore segregation is not likely to occur, cf. Section 4.

Overall, we shed light on the influence of the structure of the underlying graph and discovered the significant impact of the community structure as an important factor on the obtained segregation strength. We reveal for random geometric graphs that already after one round a provable tendency is apparent and a strong segregation occurs.

2 Model and Preliminaries

Let $G = (V, E)$ be an unweighted and undirected graph, with vertex set V and edge set E . For any vertex $v \in V$, we denote the *neighborhood* of v in G by $N_v = \{u \in V : \{u, v\} \in E\}$ and the degree of v in G by $\delta_v = |N_v|$. We consider *random geometric graphs* and *Erdős–Rényi graphs* with a total of $n \in \mathbf{N}^+$ vertices and an *expected average degree* $\bar{\delta} > 0$.

For a given $r \in \mathbf{R}^+$, a random geometric graph $G \sim \mathcal{G}(n, r)$ is obtained by distributing n vertices uniformly at random in some geometric ground space and connecting vertices u and v if and only if $\text{dist}(u, v) \leq r$. We use a two-dimensional toroidal Euclidean space with total area 1 as ground space. More formally, each vertex v is assigned to a point $(v_1, v_2) \in [0, 1]^2$ and the distance between $u = (u_1, u_2)$ and v is $\text{dist}(u, v) = \sqrt{|u_1 - v_1|_0^2 + |u_2 - v_2|_0^2}$ for $|u_i - v_i|_0 = \min\{|u_i - v_i|, 1 - |u_i - v_i|\}$. We note that using a torus instead of, e.g., a unit square, has the advantage that we do not have to consider edge cases, for vertices that are close to the boundary. In fact, a disk of radius r around any point has the same area πr^2 . Since we consider a ground space with total area 1, $r \leq \frac{1}{\sqrt{\pi}}$. As every vertex v is connected to all vertices in the disk of radius r around it, its expected average degree is $\bar{\delta} = (n-1)\pi r^2$.

For a given $p \in [0, 1]$, let $\mathcal{G}(n, p)$ denote an Erdős–Rényi graph. Each edge $\{u, v\}$ is included with probability p , independently from every other edge. It holds that $\bar{\delta} = (n-1)p$.

Consider two different vertices u and v . Let $N_{u \cap v} := |N_u \cap N_v|$ be the number of vertices in the *common neighborhood*, let $N_{u \setminus v} := |N_u \setminus N_v|$ be the number of vertices in the *exclusive neighborhood* of u , and let $N_{v \setminus u} := |N_v \setminus N_u|$ be the number of vertices in the exclusive neighborhood of v . Furthermore, with $N_{\overline{u \cup v}} := |V \setminus (N_u \cup N_v)|$, we denote the number of vertices that are neither adjacent to u nor to v .

Let G be a graph where each vertex represents an agent of type t^+ or t^- . The type of each agent is chosen independently and uniformly at random. An edge $\{u, v\}$ is *monochrome* if and only if u and v are of the same type. The *Flip Schelling Process* (FSP) is defined as follows: an agent v whose type is aligned with the type of more than $\delta_v/2$ of its neighbors keeps its type. If more than $\delta_v/2$ neighbors have a different type, then agent v changes its type. In case of a tie, i.e., if exactly $\delta_v/2$ neighbors have a different type, then v changes its type with probability $\frac{1}{2}$. FSP is a simultaneous-move, one-shot process, i.e., all agents make their decision at the same time and, moreover, only once.

For $x, y \in \mathbf{N}$, we define $[x..y] = [x, y] \cap \mathbf{N}$ and for $x \in \mathbf{N}^+$, we define $[x] = [1..x]$.

2.1 Useful Technial Lemmas

In this section, we state several lemmas that we will use in order to prove our results in the next sections.

► **Lemma 1.** *Let $X \sim \text{Bin}(n, p)$ and $Y \sim \text{Bin}(n, q)$ with $p \geq q$ be independent. Then $\Pr[X \geq Y] \geq \frac{1}{2}$.*

Proof. Let Y_1, \dots, Y_n be the individual Bernoulli trials for Y , i.e., $Y = \sum_{i \in [n]} Y_i$. Define new random variables Y'_1, \dots, Y'_n such that $Y_i = 1$ implies $Y'_i = 1$ and if $Y_i = 0$, then $Y'_i = 1$ with probability $(p - q)/(1 - q)$ and $Y'_i = 0$ otherwise. Note that for each individual Y'_i , we have $Y'_i = 1$ with probability p , i.e., $Y' = \sum_{i \in [n]} Y'_i \sim \text{Bin}(n, p)$. Moreover, as $Y' \geq Y$ for every outcome, we have $\Pr[X \geq Y] \geq \Pr[X \geq Y']$. It remains to show that $\Pr[X \geq Y'] \geq \frac{1}{2}$.

As X and Y' are equally distributed, we have $\Pr[X \geq Y'] = \Pr[X \leq Y']$. Moreover, as one of the two inequalities holds in any event, we get $\Pr[X \geq Y'] + \Pr[X \leq Y'] \geq 1$, and thus equivalently $2\Pr[X \geq Y'] \geq 1$, which proves the claim. ◀

► **Lemma 2.** *Let $n \in \mathbf{N}^+$, $p \in [0, 1)$, and let $X \sim \text{Bin}(n, p)$. Then, for all $i \in [0..n]$, it holds that $\Pr[X = i] \leq \Pr[X = \lfloor p(n + 1) \rfloor]$.*

Proof. We interpret the distribution of X as a finite series and consider the sign of the differences of two neighboring terms. A maximum of the distribution of X is located at the position at which the difference switches from positive to negative. To this end, let $b: [0, n - 1] \rightarrow [-1, 1]$ be defined such that, for all $i \in [0, n - 1] \cap \mathbf{N}$, it holds that

$$b(d) = \binom{n}{d+1} p^{d+1} (1-p)^{n-d-1} - \binom{n}{d} p^d (1-p)^{n-d}.$$

We are interested in the sign of b . In more detail, for any $d \in [0, n - 2] \cap \mathbf{N}$, if $\text{sgn}(b(d)) \geq 0$ and $\text{sgn}(b(d + 1)) \leq 0$, then $d + 1$ is a local maximum. If the sign is always negative, then there is a global maximum in the distribution of X at position 0.

In order to determine the sign of b , for all $i \in [0..n - 1]$, we rewrite

$$\begin{aligned} b(i) &= \frac{n!}{i!(n-i-1)!} p^i (1-p)^{n-i-1} \frac{p}{i+1} - \frac{n!}{i!(n-i-1)!} p^d (1-p)^{n-i-1} \frac{1-p}{n-i} \\ &= \frac{n!}{i!(n-i-1)!} p^i (1-p)^{n-i-1} \left(\frac{p}{i+1} - \frac{1-p}{n-i} \right). \end{aligned}$$

Since the term $n! p^i (1-p)^{n-i-1}$ is always non-negative, the sign of $b(i)$ is determined by the sign of $p/(i + 1) - (1 - p)/(n - i)$.

Solving for i , we get that

$$\frac{p}{i+1} - \frac{1-p}{n-i} \geq 0 \Leftrightarrow i \leq p(n+1) - 1.$$

Note that $p(n + 1) - 1$ may not be integer. Further note that the distribution of X is unimodal, as the sign of b changes at most once. Thus, each local maximum is also a global maximum. As discussed above, the largest value $d \in [0, n - 2] \cap \mathbf{N}$ such that $\text{sgn}(b(d)) \geq 0$ and $\text{sgn}(b(d + 1)) \leq 0$ then results in a global maximum at position $d + 1$. Since d needs to be integer, the largest value that satisfies this constraint is $\lfloor p(n + 1) - 1 \rfloor$. If the sign of b is always negative ($p \leq 1/(n + 1)$), then the distribution of X has a global maximum at 0, which is also satisfied by $\lfloor p(n + 1) - 1 \rfloor + 1$, which concludes the proof. ◀

► **Theorem 3** (Stirling's Formula [24, page 54]). *For all $n \in \mathbf{N}^+$, it holds that*

$$\sqrt{2\pi n}^{n+1/2} e^{-n} \cdot e^{(12n+1)^{-1}} < n! < \sqrt{2\pi n}^{n+1/2} e^{-n} \cdot e^{(12n)^{-1}}.$$

► **Corollary 4.** *For all $n \geq 2$ with $n \in \mathbf{N}$, it holds that*

$$n! > \sqrt{2\pi n}^{n+1/2} e^{-n} \quad \text{and} \tag{1}$$

$$n! < e n^{n+1/2} e^{-n}. \tag{2}$$

Proof. For both inequalities, we aim at using Theorem 3.

Equation (1): Note that $e^{(12n+1)^{-1}} > 1$, since $\frac{1}{12n+1} > 0$. Hence,

$$\sqrt{2\pi n}^{n+1/2} e^{-n} < \sqrt{2\pi n}^{n+1/2} e^{-n} \cdot e^{(12n+1)^{-1}}.$$

Equation (2): We prove this case by showing that

$$\sqrt{2\pi} e^{(12n)^{-1}} < e. \tag{3}$$

29:6 The Flip Schelling Process on Random Graphs

Note, that $e^{(12n)^{-1}}$ is strictly decreasing. Hence, we only have to check whether Equation (3) holds for $n = 2$.

$$\sqrt{2\pi}e^{(12n)^{-1}} \leq \sqrt{2\pi}e^{\frac{1}{24}} < 2.7 < e. \quad \blacktriangleleft$$

► **Lemma 5.** *Let A , B , and C be random variables such that $\Pr[A > C \wedge B > C] > 0$ and $\Pr[A > C \wedge B \leq C] > 0$. Then $\Pr[A > B \wedge A > C] \geq \Pr[A > B] \cdot \Pr[A > C]$.*

Proof. Using the definition of conditional probability, we obtain

$$\Pr[A > B \wedge A > C] = \Pr[A > B \mid A > C] \cdot \Pr[A > C].$$

Hence, we are left with bounding $\Pr[A > B \mid A > C] \geq \Pr[A > B]$. Partitioning the sample space into the two events $B > C$ and $B \leq C$ and using the law of total probability, we obtain

$$\begin{aligned} \Pr[A > B \mid A > C] &= \Pr[B > C \mid A > C] \cdot \Pr[A > B \mid A > C \wedge B > C] \\ &\quad + \Pr[B \leq C \mid A > C] \cdot \Pr[A > B \mid A > C \wedge B \leq C]. \end{aligned}$$

Note that the condition $A > C \wedge B \leq C$ already implies $A > B$ and thus the last probability equals to 1. Moreover, using the definition of conditional probability, we obtain

$$\begin{aligned} \Pr[A > B \mid A > C] &= \Pr[B > C \mid A > C] \cdot \frac{\Pr[A > B \wedge A > C \wedge B > C]}{\Pr[A > C \wedge B > C]} \\ &\quad + \Pr[B \leq C \mid A > C]. \end{aligned}$$

Using that $\Pr[B > C \mid A > C] \geq \Pr[A > C \wedge B > C]$, that $A > B \wedge B > C$ already implies $A > C$, that $\Pr[B \leq C \mid A > C] \geq \Pr[A > B \wedge B \leq C]$, and finally the law of total probability, we obtain

$$\begin{aligned} \Pr[A > B \mid A > C] &\geq \Pr[A > B \wedge A > C \wedge B > C] + \Pr[B \leq C \mid A > C] \\ &= \Pr[A > B \wedge B > C] + \Pr[B \leq C \mid A > C] \\ &\geq \Pr[A > B \wedge B > C] + \Pr[A > B \wedge B \leq C] \\ &= \Pr[A > B]. \quad \blacktriangleleft \end{aligned}$$

3 Monochrome Edges in Geometric Random Graphs

In this section, we prove the following main theorem.

► **Theorem 6.** *Let $G \sim \mathcal{G}(n, r)$ be a random geometric graph with expected average degree $\bar{\delta} = o(\sqrt{n})$. The expected fraction of monochrome edges after the FSP is at least*

$$\frac{1}{2} + \frac{9}{800} \cdot \left(\frac{1}{2} - \frac{1}{\sqrt{2\pi \lfloor \bar{\delta}/2 \rfloor}} \right)^2 \cdot \left(1 - e^{-\bar{\delta}/2} \left(1 + \frac{\bar{\delta}}{2} \right) \right) \cdot (1 - o(1)).$$

Note that the bound in Theorem 6 is bounded away from $\frac{1}{2}$ for all $\bar{\delta} \geq 2$. Moreover, the two factors depending on $\bar{\delta}$ go to $\frac{1}{2}$ and 1, respectively, for a growing $\bar{\delta}$.

Given an edge $\{u, v\}$, we prove the above lower bound on the probability that $\{u, v\}$ is monochrome in the following four steps.

1. For a vertex set, we introduce the concept of *decisiveness* that measures how much the majority is ahead of the minority in the FSP. With this, we give a lower bound on the probability that $\{u, v\}$ is monochrome based on the probability that the common neighborhood of u and v is more decisive than their exclusive neighborhoods.

2. We show that large neighborhoods are likely to be more decisive than small neighborhoods. To this end, we give bounds on the likelihood that two similar random walks behave differently. This step reduces the question of whether the common neighborhood is more decisive than the exclusive neighborhoods to whether the former is larger than the latter.
3. Turning to geometric random graphs, we show that the common neighborhood is sufficiently likely to be larger than the exclusive neighborhoods if the geometric region corresponding to the former is sufficiently large. We do this by first showing that the actual distribution of the neighborhood sizes is well approximated by independent binomial random variables. Then, we give the desired bounds for these random variables.
4. We show that the existence of the edge $\{u, v\}$ in the geometric random graph makes it sufficiently likely that the geometric region hosting the common neighborhood of u and v is sufficiently large.

3.1 Monochrome Edges via Decisive Neighborhoods

Let $\{u, v\}$ be an edge of a given graph. To formally define the above mentioned decisiveness, let $N_{u \cap v}^+$ and $N_{u \cap v}^-$ be the number of vertices in the common neighborhood of u and v that are occupied by agents of type t^+ and t^- , respectively. Then $D_{u \cap v} := |N_{u \cap v}^+ - N_{u \cap v}^-|$ is the *decisiveness* of the common neighborhood of u and v . Analogously, we define $D_{u \setminus v}$ and $D_{v \setminus u}$ for the exclusive neighborhoods of u and v , respectively.

The following theorem bounds the probability for $\{u, v\}$ to be monochrome based on the probability that the common neighborhood is more decisive than each of the exclusive ones.

► **Theorem 7.** *In the FSP, let $\{u, v\} \in E$ be an edge and let D be the event $\{D_{u \cap v} > D_{u \setminus v} \wedge D_{u \cap v} > D_{v \setminus u}\}$. Then $\{u, v\}$ is monochrome with probability at least $1/2 + \Pr[D]/2$.*

Proof. If D occurs, then the types of u and v after the FSP coincide with the predominant type before the FSP in the shared neighborhood. Thus, $\{u, v\}$ is monochrome.

Otherwise, assuming \bar{D} , w.l.o.g., let $D_{u \cap v} \leq D_{u \setminus v}$ and assume further the type of v has already been determined. If $D_{u \cap v} = D_{u \setminus v}$, then u chooses a type uniformly at random, which coincides with the type of v with probability $\frac{1}{2}$. Otherwise, $D_{u \cap v} < D_{u \setminus v}$ and thus u takes the type that is predominant in u 's exclusive neighborhood, which is t^+ and t^- with probability $\frac{1}{2}$, each. Moreover, this is independent from the type of v as v 's neighborhood is disjoint to u 's exclusive neighborhood.

Thus, for the event M that $\{u, v\}$ is monochrome, we get $\Pr[M | D] = 1$ and $\Pr[M | \bar{D}] = \frac{1}{2}$. Hence, we get $\Pr[M] > \Pr[D] + \frac{1}{2}(1 - \Pr[D]) = \frac{1}{2} + \Pr[D]/2$. ◀

3.2 Large Neighborhoods are More Decisive

The goal of this section is to reduce the question of how decisive a neighborhood is to the question of how large it is. To be more precise, assume we have a set of vertices of size a and give each vertex the type t^+ and t^- , respectively, each with probability $\frac{1}{2}$. Let A_i for $i \in [a]$ be the random variable that takes the value $+1$ and -1 if the i -th vertex in this set has type t^+ and t^- , respectively. Then, for $A = \sum_{i \in [a]} A_i$, the decisiveness of the vertex set is $|A|$. In the following, we study the decisiveness $|A|$ depending on the size a of the set. Note that this can be viewed as a random walk on the integer line: Starting at 0, in each step, it moves one unit either to the left or to the right with equal probabilities. We are interested in the distance from 0 after a steps.

Assume for the vertices u and v that we know that b vertices lie in the common neighborhood and a vertices lie in the exclusive neighborhood of u . Moreover, let A and B be the positions of the above random walk after a and b steps, respectively. Then the event

29:8 The Flip Schelling Process on Random Graphs

$D_{u \cap v} > D_{u \setminus v}$ is equivalent to $|B| > |A|$. Motivated by this, we study the probability of $|B| > |A|$, assuming $b \geq a$. The core difficulty here comes from the fact that we require $|B|$ to be strictly larger than $|A|$. Also note that $a + b$ corresponds to the degree of u in the graph. Thus, we have to study the random walks also for small numbers of a and b . We note that all results in this section are independent from the specific application to the FSP, and thus might be of independent interest.

Before we give a lower bound on the probability that $|B| > |A|$, we need the following technical lemma. It states that doing more steps in the random walk only makes it more likely to deviate further from the starting position.

► **Lemma 8.** *For $i \in [a]$ and $j \in [b]$ with $0 \leq a \leq b$, let A_i and B_j be independent random variables that are -1 and 1 each with probability $\frac{1}{2}$. Let $A = \sum_{i \in [a]} A_i$ and $B = \sum_{j \in [b]} B_j$. Then $\Pr[|A| < |B|] \geq \Pr[|A| > |B|]$.*

Proof. Let Δ_k be the event that $|B| - |A| = k$. First note that

$$\Pr[|A| < |B|] = \sum_{k \in [b]} \Pr[\Delta_k] \quad \text{and} \quad \Pr[|A| > |B|] = \sum_{k \in [a]} \Pr[\Delta_{-k}].$$

To prove the statement of the lemma, it thus suffices to prove the following claim.

▷ **Claim 9.** For $k \geq 0$, $\Pr[\Delta_k] \geq \Pr[\Delta_{-k}]$.

We prove this claim via induction on $b - a$. For the base case $a = b$, A and B are equally distributed and thus Claim 9 clearly holds.

For the induction step, let B^+ be the random variable that takes the values $B + 1$ and $B - 1$ with probability $\frac{1}{2}$ each. Note that B^+ represents the same type of random walk as A and B but with $b + 1$ steps. Moreover B^+ is coupled with B to make the same decisions in the first b steps. Let Δ_k^+ be the event that $|B^+| - |A| = k$. It remains to show that Claim 9 holds for these Δ_k^+ . For this, first note that the claim trivially holds for $k = 0$. For $k \geq 1$, we can use the definition of Δ_k^+ and the induction hypothesis to obtain

$$\begin{aligned} \Pr[\Delta_k^+] &= \frac{\Pr[\Delta_{k-1}]}{2} + \frac{\Pr[\Delta_{k+1}]}{2} \\ &\geq \frac{\Pr[\Delta_{-k+1}]}{2} + \frac{\Pr[\Delta_{-k-1}]}{2} = \Pr[\Delta_{-k}^+]. \end{aligned} \quad \blacktriangleleft$$

Using Lemma 8, we now prove the following general bound for the probability that $|A| < |B|$, depending on certain probabilities for binomially distributed variables.

► **Lemma 10.** *For $i \in [a]$ and $j \in [b]$ with $0 \leq a \leq b$, let A_i and B_j be independent random variables that are -1 and 1 each with probability $\frac{1}{2}$. Let $A = \sum_{i \in [a]} A_i$ and $B = \sum_{j \in [b]} B_j$. Moreover, let $X \sim \text{Bin}(a, \frac{1}{2})$, $Y \sim \text{Bin}(b, \frac{1}{2})$, and $Z \sim \text{Bin}(a + b, \frac{1}{2})$. Then*

$$\Pr[|A| < |B|] \geq \frac{1}{2} - \Pr\left[Z = \frac{a+b}{2}\right] + \frac{\Pr[X = \frac{a}{2}] \cdot \Pr[Y = \frac{b}{2}]}{2}.$$

Proof. Using that $\Pr[|A| < |B|] \geq \Pr[|A| > |B|]$ (see Lemma 8), we obtain

$$\begin{aligned} &\Pr[|A| < |B|] + \Pr[|A| > |B|] + \Pr[|A| = |B|] = 1 \\ \Rightarrow &2\Pr[|A| < |B|] + \Pr[|A| = |B|] \geq 1 \\ \Leftrightarrow &\Pr[|A| < |B|] \geq \frac{1}{2} - \frac{\Pr[|A| = |B|]}{2}. \end{aligned} \quad (4)$$

Thus, it remains to give an upper bound for $\Pr[|A| = |B|]$.

Using the inclusion–exclusion principle and the fact that B is symmetric around 0, i.e., $\Pr[B = x] = \Pr[B = -x]$ for any x , we obtain

$$\begin{aligned} \Pr[|A| = |B|] &= \Pr[A = B \vee A = -B] \\ &= \Pr[A = B] + \Pr[A = -B] - \Pr[A = B = 0] \\ &= 2\Pr[A = -B] - \Pr[A = B = 0]. \end{aligned} \tag{5}$$

We estimate $\Pr[A = -B]$ and $\Pr[A = B = 0]$ using bounds for binomially distributed variables. To this end, define new random variables $X_i = \frac{A_i+1}{2}$ for $i \in [a]$ and let $X = \sum_{i \in [a]} X_i$. Note that the X_i are independent and take values 0 and 1, each with probability $\frac{1}{2}$. Thus, $X \sim \text{Bin}(a, \frac{1}{2})$. Moreover, $A = 2X - a$. Analogously, we define Y with $Y \sim \text{Bin}(b, \frac{1}{2})$ and $B = 2Y - b$. Note that X and Y are independent and thus $Z = X + Y \sim \text{Bin}(a + b, \frac{1}{2})$. With this, we get

$$\begin{aligned} \Pr[A = -B] &= \Pr[2X - a = -2Y + b] = \Pr\left[Z = \frac{a+b}{2}\right], \text{ and} \\ \Pr[A = B = 0] &= \Pr[A = 0] \cdot \Pr[B = 0] = \Pr\left[X = \frac{a}{2}\right] \cdot \Pr\left[Y = \frac{b}{2}\right]. \end{aligned}$$

This, together with Equations (4) and (5) yield the claim. ◀

The bound in Lemma 10 becomes worse for smaller values of a and b . Considering this worst case, we obtain the following specific bound.

► **Theorem 11.** *For $i \in [a]$ and $j \in [b]$ with $0 \leq a \leq b$, let A_i and B_j be independent random variables that are -1 and 1 each with probability $\frac{1}{2}$. Let $A = \sum_{i \in [a]} A_i$ and $B = \sum_{j \in [b]} B_j$. If $a = b = 0$ or $a = b = 1$, then $\Pr[|A| < |B|] = 0$. Otherwise $\Pr[|A| < |B|] \geq \frac{3}{16}$.*

Proof. Clearly, if $a = b = 0$, then $A = B = 0$ and thus $\Pr[|A| < |B|] = 0$. Similarly, if $a = b = 1$, then $|A| = |B| = 1$ and thus $\Pr[|A| < |B|] = 0$. For the remainder, assume that neither $a = b = 0$ nor $a = b = 1$, and let X, Y , and Z be defined as in Lemma 10, i.e., $X \sim \text{Bin}(a, \frac{1}{2})$, $Y \sim \text{Bin}(b, \frac{1}{2})$, and $Z \sim \text{Bin}(a + b, \frac{1}{2})$.

If $a + b$ is odd, then $\Pr[Z = \frac{a+b}{2}] = 0$. Thus, by Lemma 10, we have $\Pr[|A| < |B|] \geq \frac{1}{2}$. If $a + b$ is even and $a + b \geq 6$, then

$$\Pr\left[Z = \frac{a+b}{2}\right] = \binom{a+b}{\frac{a+b}{2}} \left(\frac{1}{2}\right)^{a+b} \leq \binom{6}{3} \left(\frac{1}{2}\right)^6 = \frac{5}{16}.$$

Hence, by Lemma 10, we have $\Pr[|A| < |B|] \geq \frac{1}{2} - \frac{5}{16} = \frac{3}{16}$.

If $a + b < 6$ (and $a + b$ is even), there are four cases: $a = 0, b = 2$; $a = 0, b = 4$; $a = 1, b = 3$; $a = 2, b = 2$. If $a = 0$ and $b = 2$, then $A = 0$ with probability 1 and $|B| = 2$ with probability $\frac{1}{2}$. Thus, $\Pr[|A| < |B|] = \frac{1}{2}$. If $a = 0$ and $b = 4$, then $|A| < |B|$ unless $B = 0$. As $\Pr[B = 0] = \binom{4}{2} \cdot (\frac{1}{2})^4 = \frac{3}{8}$, we get $\Pr[|A| < |B|] = 1 - \frac{3}{8} = \frac{5}{8}$. If $a = 1$ and $b = 3$, then $|A| = 1$ with probability 1 and $|B| = 3$ with probability $\frac{1}{4}$ (either $B_1 = B_2 = B_3 = 1$ or $B_1 = B_2 = B_3 = -1$). Thus, $\Pr[|A| < |B|] = \frac{1}{4}$. If $a = b = 2$, then $|A| = 0$ with probability $\frac{1}{2}$ and $|B| = 2$ with probability $\frac{1}{2}$. Thus $\Pr[|A| < |B|] = \frac{1}{4}$.

We note that the bound of $\Pr[|A| < |B|] = \frac{3}{16}$ is tight for $a = b = 3$. ◀

3.3 Large Common Regions Yield Large Common Neighborhoods

To be able to apply Theorem 11 to an edge $\{u, v\}$, we need to make sure that the size of their common neighborhood (corresponding to b in the theorem) is at least the size of the exclusive neighborhoods (corresponding to a in the theorem). In the following, we give bounds for the probability that this happens. Note that this is the first time we actually take the graph into account. Thus, all above considerations hold for arbitrary graphs.

Recall that we consider random geometric graphs $\mathcal{G}(n, r)$ and u and v are each connected to all vertices that lie within a disk of radius r around them. As u and v are adjacent, their disks intersect, which separates the ground space into four regions; cf. Fig 2a. Let $R_{u \cap v}$ be the intersection of the two disks. Let $R_{u \setminus v}$ be the set of points that lie in the disk of u but not in the disk of v , and analogously, let $R_{v \setminus u}$ be the disk of v minus the disk of u . Finally, let $R_{\overline{u \cup v}}$ be the set of points outside both disks. Then, each of the $n - 2$ remaining vertices ends up in exactly one of these regions with a probability equal to the corresponding measure. Let $\mu(\cdot)$ be the area of the respective region and $p = \mu(R_{u \cap v})$ and $q = \mu(R_{u \setminus v}) = \mu(R_{v \setminus u})$ be the probabilities for a vertex to lie in the common and exclusive regions, respectively. The probability for $R_{\overline{u \cup v}}$ is then $1 - p - 2q$.

We are now interested in the sizes $N_{u \cap v}$, $N_{u \setminus v}$, and $N_{v \setminus u}$ of the common and the exclusive neighborhoods, respectively. As each of the $n - 2$ remaining vertices ends up in $N_{u \cap v}$ with probability p , we have $N_{u \cap v} \sim \text{Bin}(n - 2, p)$. For $N_{u \setminus v}$ and $N_{v \setminus u}$, we already know that v is a neighbor of u and vice versa. Thus, $(N_{u \setminus v} - 1) \sim \text{Bin}(n - 2, q)$ and $(N_{v \setminus u} - 1) \sim \text{Bin}(n - 2, q)$. Moreover, the three random variables are not independent, as each vertex lies in only exactly one of the four neighborhoods, i.e., $N_{u \cap v}$, $(N_{u \setminus v} - 1)$, $(N_{v \setminus u} - 1)$, and the number of vertices in neither neighborhood together follow a multinomial distribution $\text{Multi}(n - 2, \mathbf{p})$ with $\mathbf{p} = (p, q, q, 1 - p - 2q)$.

The following lemma shows that these dependencies are small if p and q are sufficiently small. This lets us assume that $N_{u \cap v}$, $(N_{u \setminus v} - 1)$, $(N_{v \setminus u} - 1)$ are independent random variables following binomial distributions if the expected average degree is not too large.

► **Lemma 12.** *Let $X = (X_1, X_2, X_3, X_4) \sim \text{Multi}(n, \mathbf{p})$ with $\mathbf{p} = (p, q, q, 1 - p - 2q)$. Then there exist independent random variables $Y_1 \sim \text{Bin}(n, p)$, $Y_2 \sim \text{Bin}(n, q)$, and $Y_3 \sim \text{Bin}(n, q)$ such that $\Pr[(X_1, X_2, X_3) = (Y_1, Y_2, Y_3)] \geq 1 - 3n \cdot \max(p, q)^2$.*

Proof. Let $Y_1 \sim \text{Bin}(n, p)$, and $Y_2, Y_3 \sim \text{Bin}(n, q)$ be independent random variables. We define the event B to hold, if each of the n individual trials increments at most one of the random variables Y_1, Y_2 , or Y_3 . More formally, for $i \in [3]$ and $j \in [n]$, let $Y_{i,j}$ be the individual Bernoulli trials of Y_i , i.e., $Y_i = \sum_{j \in [n]} Y_{i,j}$. For $j \in [n]$, we define the event B_j to be $Y_{1,j} + Y_{2,j} + Y_{3,j} \leq 1$, and the event $B = \bigcap_{j \in [n]} B_j$.

Based on this, we now define the random variables X_1, X_2, X_3 , and X_4 as follows. If B holds, we set $X_i = Y_i$ for $i \in [3]$ and $X_4 = n - X_1 - X_2 - X_3$. Otherwise, if \overline{B} , we draw $X = (X_1, X_2, X_3, X_4) \sim \text{Multi}(n, \mathbf{p})$ independently from Y_1, Y_2 , and Y_3 with $\mathbf{p} = (p, q, q, 1 - p - 2q)$. Note that X clearly follows $\text{Multi}(n, \mathbf{p})$ if \overline{B} . Moreover, conditioned on B , each individual trial increments exactly one of the variables X_1, X_2, X_3 , or X_4 with probabilities p, q, q , and $1 - p - 2q$, respectively, i.e., $X \sim \text{Multi}(n, \mathbf{p})$.

Thus, we end up with $X \sim \text{Multi}(n, \mathbf{p})$. Additionally, we have three independent random variables $Y_1 \sim \text{Bin}(n, p)$, and $Y_2, Y_3 \sim \text{Bin}(n, q)$ with $(X_1, X_2, X_3) = (Y_1, Y_2, Y_3)$ if B holds. Thus, to prove the lemma, it remains to show that $\Pr[B] \geq 1 - 3n \max(p, q)^2$. For $j \in [n]$, the probability that the j th trial goes wrong is

$$\begin{aligned}\Pr[\overline{B}_j] &= 1 - ((1-p)(1-q)^2) - (p(1-q)^2) - 2(q(1-p)(1-q)) \\ &= 2pq - 2pq^2 + q^2 \leq 2pq + q^2 \leq 3 \cdot \max(p, q)^2.\end{aligned}$$

Using the union bound it follows that $\Pr[\overline{B}] \leq \sum_{j \in [n]} \Pr[\overline{B}_j] \leq 3n \cdot \max(p, q)^2$. ◀

As mentioned before, we are interested in the event $N_{u \cap v} \geq N_{u \setminus v}$ (and likewise $N_{u \cap v} \geq N_{v \setminus u}$), in order to apply Theorem 11. Moreover, due to Lemma 12, we know that $N_{u \cap v}$ and $(N_{u \setminus v} - 1)$ almost behave like independent random variables that follow $\text{Bin}(n-2, p)$ and $\text{Bin}(n-2, q)$, respectively. The following lemma helps to bound the probability for $N_{u \cap v} \geq N_{u \setminus v}$. Note that it gives a bound for the probability of achieving strict inequality (instead of just \geq), which accounts for the fact that $(N_{u \setminus v} - 1)$ and not $N_{u \setminus v}$ itself follows a binomial distribution.

► **Lemma 13.** *Let $n \in \mathbf{N}$ with $n \geq 2$, and let $p \geq q > 0$. Further, let $X \sim \text{Bin}(n, p)$ and $Y \sim \text{Bin}(n, q)$ be independent, let $d = \lfloor p(n+1) \rfloor$, and assume $d = o(\sqrt{n})$, then $\Pr[X > Y] \geq (\frac{1}{2} - 1/\sqrt{2\pi d})(1 - o(1))$.*

Proof. By Lemma 1, we get $\Pr[X \geq Y] \geq \frac{1}{2}$, and we bound

$$\Pr[X > Y] = \Pr[X \geq Y] - \Pr[X = Y] \geq \frac{1}{2} - \Pr[X = Y],$$

leaving us to bound $\Pr[X = Y]$ from above. By independence of X and Y , we get

$$\Pr[X = Y] = \sum_{i \in [n]} \Pr[X = i] \cdot \Pr[Y = i]. \quad (6)$$

Note that, by Lemma 2, for all $i \in [0..n]$, it holds that $\Pr[X = i] \leq \Pr[X = d]$. Assume that we have a bound B such that $\Pr[X = d] \leq B$. Substituting this into Equation (6) yields

$$\Pr[X = Y] \leq B \sum_{i \in [n]} \Pr[Y = i] = B,$$


resulting in $\Pr[X > Y] \geq \frac{1}{2} - B$. Thus, we now derive such a bound for B and apply the inequality that for all $x \in \mathbf{R}$, it holds that $1 + x \leq e^x$, as well as Equation (1). We get

$$\begin{aligned}\binom{n}{d} p^d (1-p)^{n-d} &\leq \frac{n^d}{d!} \left(\frac{d}{n}\right)^d \left(1 - \frac{d}{n}\right)^n \left(1 - \frac{d}{n}\right)^{-d} \\ &\leq \frac{d^d}{d!} e^{-d} \left(1 - \frac{d}{n}\right)^{-d} \\ &\leq \frac{d^d}{\sqrt{2\pi d}^{d+1/2} e^{-d}} e^{-d} \left(1 - \frac{d}{n}\right)^{-d} \\ &= \frac{1}{\sqrt{2\pi d}} \frac{1}{(1 - d/n)^d}.\end{aligned} \quad (7)$$

By Bernoulli's inequality, we bound $(1 - d/n)^d \geq 1 - d^2/n = 1 - o(1)$ by the assumption $d = o(\sqrt{n})$. Substituting this back into Equation (7) concludes the proof. ◀

Finally, in order to apply Theorem 11, we have to make sure not to end up in the special case where $a = b \leq 1$, i.e., we have to make sure that the common neighborhood includes at least two vertices. The probability for this to happen is given by the following lemma.

29:12 The Flip Schelling Process on Random Graphs



(a) The geometric regions corresponding to the common and exclusive neighborhoods, respectively, with yellow illustrating $R_{u \cap v}$ and blue illustrating $R_{u \setminus v}$ and $R_{v \setminus u}$.

(b) Let α be the central angle determined by the intersection points i and j , and let x be the corresponding circular sector (illustrated in yellow).

(c) Let y be a triangle in the intersection (illustrated in green) determined by the radical axis ℓ and the central angle α , cf. Fig 2b.

(d) The height h divides the area $\mu(y)$ (illustrated in green) of the triangle y , cf. Fig 2c, into two subareas of equal size, since adjacent and opposite legs have the same length r .

■ **Figure 2** The neighborhood of two adjacent vertices u and v in a random geometric graph.

► **Lemma 14.** Let $X \sim \text{Bin}(n, p)$ and let $c = np \in o(n)$. Then it holds that $\Pr[X > 1] \geq (1 - e^{-c}(1 + c))(1 - o(1))$.

Proof. As $X > 1$ holds if and only if $X \neq 0$ and $X \neq 1$, we get

$$\Pr[X > 1] = 1 - \Pr[X = 0] - \Pr[X = 1] = 1 - (1 - p)^n - n \cdot p \cdot (1 - p)^{n-1}.$$

Using that for all $x \in \mathbf{R}$ it holds that $1 - x \leq e^{-x}$, we get

$$\begin{aligned} \Pr[X > 1] &\geq 1 - e^{-pn} - n \cdot p \cdot e^{-p(n-1)} \\ &= 1 - e^{-c} - c \cdot e^{c/n} \cdot e^{-c} \\ &= 1 - e^{-c} \left(1 + c \cdot e^{c/n}\right). \end{aligned}$$

As $e^{c/n}$ goes to 1 for $n \rightarrow \infty$, we get the claimed bound. ◀

3.4 Many Edges Have Large Common Regions

In Section 3.3, we derived a lower bound on the probability that $N_{u \cap v} \geq N_{u \setminus v}$ provided that the probability for a vertex to end up in the shared region $R_{u \cap v}$ is sufficiently large compared to $R_{u \setminus v}$. In the following, we estimate the measures of these regions depending on the distance between u and v . Then, we give a lower bound on the probability that $\mu(R_{u \cap v}) \geq \mu(R_{u \setminus v})$.

► **Lemma 15.** Let $G \sim \mathcal{G}(n, r)$ be a random geometric graph with expected average degree $\bar{\delta}$, let $\{u, v\} \in E$ be an edge, and let $\tau := \frac{\text{dist}(u, v)}{r}$. Then,

$$\mu(R_{u \cap v}) = \frac{\bar{\delta}}{(n-1)\pi} \left(2 \arccos\left(\frac{\tau}{2}\right) - \sin\left(2 \arccos\left(\frac{\tau}{2}\right)\right) \right) \text{ and} \quad (8)$$

$$\mu(R_{u \setminus v}) = \mu(R_{v \setminus u}) = \frac{\bar{\delta}}{n-1} - \mu(R_{u \cap v}). \quad (9)$$

Proof. We start with proving Equation (8). Let i and j be the two intersection points of the disks of u and v , let α be the central angle enclosed by i and j , and let x be the corresponding circular sector, cf. Fig 2b. Moreover, let the triangle y be a subarea of x determined by α

and the radical axis ℓ , cf. Fig 2c. Let h denote the height of the triangle y , cf. Fig 2d. For our calculations, we restrict the length of ℓ by the intersection points i and j . Since we consider the intersection between disks and thus ℓ divides the area $\mu(R_{u \cap v})$ into two subareas of equal sizes, it holds that $\mu(R_{u \cap v}) = 2(\mu(x) - \mu(y))$. Considering the two areas $\mu(x)$ and $\mu(y)$, it holds that

$$\mu(x) = \frac{\alpha}{2}r^2 \quad \text{and} \quad \mu(y) = h \cdot \frac{\ell}{2} = \cos\left(\frac{\alpha}{2}\right)r \cdot \sin\left(\frac{\alpha}{2}\right)r = \frac{\sin(\alpha)}{2}r^2. \quad (10)$$

For the central angle α we know $\cos(\alpha/2) = h/r = \tau/2$ and therefore $\alpha = 2 \arccos(\frac{\tau}{2})$. Together with Equation (10), we obtain

$$\mu(R_{u \cap v}) = 2(\mu(x) - \mu(y)) = 2 \left(\frac{2 \arccos(\frac{\tau}{2})}{2}r^2 - \frac{\sin(2 \arccos(\frac{\tau}{2}))}{2}r^2 \right). \quad (11)$$

The area of a general circle is equal to πr^2 . Since we consider a ground space with total area 1, the area of one disk in the random geometric graph equals $\frac{\delta}{n-1}$, i.e., $r^2 = \frac{\delta}{(n-1)\pi}$. Together with Equation (11), we obtain Equation (8).

Equation (9): We get the claimed equality by noting that $\mu(R_{u \cap v}) + \mu(R_{u \setminus v}) = \pi r^2$. ◀

► **Lemma 16.** *Let $G \sim \mathcal{G}(n, r)$ be a random geometric graph, and let $\{u, v\} \in E$ be an edge. Then $\Pr[\mu(R_{u \cap v}) \geq \mu(R_{u \setminus v})] \geq (\frac{4}{5})^2$.*

Proof. Let $\tau = \frac{\text{dist}(u,v)}{r}$. By Lemma 15 with $\mu(R_{u \cap v}) \geq \mu(R_{v \setminus u})$, we get

$$\left(2 \arccos\left(\frac{\tau}{2}\right) - \sin\left(2 \arccos\left(\frac{\tau}{2}\right)\right) \right) \geq \frac{\pi}{2},$$

which is true for $\tau \geq \frac{4}{5}$. The area of a disk of radius $\frac{4}{5}r$ is $(\pi(\frac{4}{5}r)^2) / (\pi r^2) = (\frac{4}{5})^2$ times the area of a disk of radius r . Hence, the fraction of edges with distance at most $\frac{4}{5}r$ is at least $(\frac{4}{5})^2$, concluding the proof. ◀

3.5 Proof of Theorem 6

By Theorem 7, the probability that a random edge $\{u, v\}$ is monochrome is at least $\frac{1}{2} + \Pr[D]/2$, where D is the event that the common neighborhood of u and v is more decisive than each exclusive neighborhood. It remains to bound $\Pr[D]$.

Existence of an edge yields a large shared region. Let R be the event that $\mu(R_{u \cap v}) \geq \mu(R_{u \setminus v})$. Note that this also implies $\mu(R_{u \cap v}) \geq \mu(R_{v \setminus u})$ as $\mu(R_{u \setminus v}) = \mu(R_{v \setminus u})$. Due to the law of total probability, we have

$$\Pr[D] \geq \Pr[R] \cdot \Pr[D | R].$$

Due to Lemma 16, we have $\Pr[R] \geq (\frac{4}{5})^2$. By conditioning on R in the following, we can assume that $\mu(R_{u \cap v}) \geq \frac{\bar{\delta}}{2n} \geq \mu(R_{u \setminus v}) = \mu(R_{v \setminus u})$, where $\bar{\delta}$ is the expected average degree.

Neighborhood sizes are roughly binomially distributed. The next step is to go from the size of the regions to the number of vertices in these regions. Each of the remaining $n' = n - 2$ vertices is sampled independently to lie in one of the regions $R_{u \cap v}$, $R_{u \setminus v}$, $R_{v \setminus u}$, or $R_{u \cup v}$. Denote the resulting numbers of vertices with X_1 , X_2 , X_3 , and X_4 , respectively. Then (X_1, X_2, X_3, X_4) follows a multinomial distribution with parameter $\mathbf{p} = (p, q, q, 1 - p - 2q)$

29:14 The Flip Schelling Process on Random Graphs

for $p = \mu(R_{u \cap v})$ and $q = \mu(R_{u \setminus v}) = \mu(R_{v \setminus u})$. Note that $N_{u \cap v} = X_1$, $N_{u \setminus v} = X_2 + 1$, and $N_{v \setminus u} = X_3 + 1$ holds for the sizes of the common and exclusive neighborhoods, where the $+1$ comes from the fact that v is always a neighbor of u and vice versa.

We apply Lemma 12 to obtain independent binomially distributed random variables Y_1 , Y_2 , and Y_3 that are likely to coincide with $X_1 = N_{u \cap v}$, $X_2 = N_{u \setminus v} - 1$, and $X_3 = N_{v \setminus u} - 1$, respectively. Let B denote the event that $(N_{u \cap v}, N_{u \setminus v} - 1, N_{v \setminus u} - 1) = (Y_1, Y_2, Y_3)$. Again, using the law of total probabilities and due to the fact that R and B are independent, we get

$$\Pr[D \mid R] \geq \Pr[B \mid R] \cdot \Pr[D \mid R \cap B] = \Pr[B] \cdot \Pr[D \mid R \cap B].$$

Note that $p, q \leq \frac{\bar{\delta}}{n}$ for the expected average degree $\bar{\delta}$. Thus, Lemma 12 implies that $\Pr[B] \geq \left(1 - 3\bar{\delta}^2/n\right)$. Conditioning on B makes it correct to assume that $N_{u \cap v} \sim \text{Bin}(n', p)$, $(N_{u \setminus v} - 1) \sim \text{Bin}(n', q)$, $(N_{v \setminus u} - 1) \sim \text{Bin}(n', q)$ are independently distributed. Additionally conditioning on R gives us $p \geq \frac{\bar{\delta}}{2n} \geq q$.

A large shared region yields a large shared neighborhood. In the next step, we consider an event that makes sure that the number $N_{u \cap v}$ of vertices in the shared neighborhood is sufficiently large. Let N_1 , N_2 , and N_3 be the events that $N_{u \cap v} \geq N_{u \setminus v}$, $N_{u \cap v} \geq N_{v \setminus u}$, and $N_{u \cap v} > 1$, respectively. Let N be the intersection of N_1 , N_2 , and N_3 . We obtain

$$\begin{aligned} \Pr[D \mid R \cap B] &\geq \Pr[N \mid R \cap B] \cdot \Pr[D \mid R \cap B \cap N] \\ &\geq \Pr[N_1 \mid R \cap B] \cdot \Pr[N_2 \mid R \cap B] \cdot \Pr[N_3 \mid R \cap B] \cdot \Pr[D \mid R \cap B \cap N], \end{aligned}$$

where the last step follows from Lemma 5 as the inequalities in N_1 , N_2 , and N_3 all go in the same direction. Note that $N_{u \cap v} \geq N_{u \setminus v}$ is equivalent to $N_{u \cap v} > N_{u \setminus v} - 1$. Due to the condition on B , $N_{u \cap v}$ and $N_{u \setminus v} - 1$ are independent random variables following $\text{Bin}(n', p)$ and $\text{Bin}(n', q)$, respectively, with $p \geq q$ due to the condition on R . Thus, we can apply Lemma 13, to obtain

$$\Pr[N_1 \mid R \cap B] = \Pr[N_2 \mid R \cap B] \geq \frac{1}{2} - \frac{1}{\sqrt{2\pi[\bar{\delta}/2](1 - o(1))}},$$

and Lemma 14 gives the bound

$$\Pr[N_3 \mid R \cap B] \geq 1 - e^{-\bar{\delta}/2} \left(1 + \frac{\bar{\delta}}{2} \cdot (1 + o(1))\right).$$

Note that both of these probabilities are bounded away from 0 for $\bar{\delta} \geq 2$. Conditioning on N lets us assume that the shared neighborhood of u and v contains at least two vertices and that it is at least as big as each of the exclusive neighborhoods.

A large shared neighborhood yields high decisiveness. The last step is to actually bound the remaining probability $\Pr[D \mid R \cap B \cap N]$. Note that, once we know the number of vertices in the shared and exclusive neighborhoods, the decisiveness no longer depends on R or B , i.e., we can bound $\Pr[D \mid N]$ instead. For this, let D_1 and D_2 be the events that $D_{u \cap v} > D_{u \setminus v}$ and $D_{u \cap v} > D_{v \setminus u}$, respectively. Note that D is their intersection. Moreover, due to Lemma 5, we have $\Pr[D \mid N] \geq \Pr[D_1 \mid N] \cdot \Pr[D_2 \mid N]$. To bound $\Pr[D_1 \mid N] = \Pr[D_2 \mid N]$, we use Theorem 11. Note that the b and a in Theorem 11 correspond to $N_{u \cap v}$ and $N_{u \setminus v} + 1$ (the $+1$ coming from the fact that $N_{u \setminus v}$ does not count the vertex v). Moreover conditioning on N implies that $a \leq b$ and $b > 1$. Thus, Theorem 11 implies $\Pr[D_1 \mid N] \geq \frac{3}{16}$.

Conclusion. The above arguments give us that the fraction of monochrome edges is

$$\frac{1}{2} + \frac{\Pr[D]}{2} \geq \frac{1}{2} + \frac{1}{2} \cdot \underbrace{\Pr[R]}_{\geq (\frac{4}{5})^2} \cdot \underbrace{\Pr[B]}_{1-o(1)} \cdot \underbrace{(\Pr[N_1 | R \cap B])^2}_{\geq \frac{1}{2} - \frac{1}{\sqrt{2\pi \lfloor \bar{\delta}/2 \rfloor}}} \cdot \underbrace{\Pr[N_3 | R \cap B]}_{\geq 1 - e^{-\bar{\delta}/2} (1 + \frac{\bar{\delta}}{2})} \cdot \underbrace{(\Pr[D_1 | N])^2}_{\geq \frac{3}{16}},$$

where we omitted the $o(1)$ terms for $\Pr[N_1 | R \cap B]$ and $\Pr[N_3 | R \cap B]$, as they are already covered by the $1 + o(1)$ coming from $\Pr[B]$. This yields the bound stated in Theorem 6:

$$\frac{1}{2} + \frac{9}{800} \cdot \left(\frac{1}{2} - \frac{1}{\sqrt{2\pi \lfloor \bar{\delta}/2 \rfloor}} \right)^2 \cdot \left(1 - e^{-\bar{\delta}/2} \left(1 + \frac{\bar{\delta}}{2} \right) \right) \cdot (1 - o(1)).$$

4 Monochrome Edges in Erdős–Rényi Graphs

In the following, we are interested in the probability that an edge $\{u, v\}$ is monochrome after the FSP on Erdős–Rényi graphs. In contrast to geometric random graphs, we prove an upper bound. To this end, we show that it is likely that the common neighborhood is empty and therefore u and v choose their types to be the predominant type in their exclusive neighborhood, which is t^+ and t^- with probability $\frac{1}{2}$, each.

► **Theorem 17.** *Let $G \sim \mathcal{G}(n, p)$ be an Erdős–Rényi graph with expected average degree $\bar{\delta} = o(\sqrt{n})$. The expected fraction of monochrome edges after the FSP is at most $\frac{1}{2} + o(1)$.*

Proof. Given an edge $\{u, v\}$, let M be the event that $\{u, v\}$ is monochrome. We first split M into disjoint sets with respect to the size of the common neighborhood and apply the law of total probability and get

$$\begin{aligned} \Pr[M] &= \Pr[M | N_{u \cap v} = 0] \cdot \Pr[N_{u \cap v} = 0] + \Pr[M | N_{u \cap v} > 0] \cdot \Pr[N_{u \cap v} > 0] \\ &\leq \Pr[M | N_{u \cap v} = 0] \cdot 1 + 1 \cdot \Pr[N_{u \cap v} > 0]. \end{aligned}$$

We bound each of the summands separately. For estimating $\Pr[M | N_{u \cap v} = 0]$, we note that the types of u and v are determined by the predominant type in disjoint vertex sets. By definition of the FSP this implies that the probability of a monochrome edge is equal to $\frac{1}{2}$.

We are left with bounding $\Pr[N_{u \cap v} > 0]$. Note that $N_{u \cap v} \sim \text{Bin}(n, p^2)$. Thus, by Bernoulli's inequality we get $\Pr[N_{u \cap v} > 0] = 1 - \Pr[N_{u \cap v} = 0] = 1 - (1 - p^2)^n \leq np^2$. Noting that $np^2 = o(1)$ holds due to our assumption on $\bar{\delta}$, concludes the proof. ◀

References

- 1 Aishwarya Agarwal, Edith Elkind, Jiarui Gan, and Alexandros A. Voudouris. Swap stability in Schelling games on graphs. In *AAAI'20*, pages 1758–1765, 2020. URL: <https://aaai.org/ojs/index.php/AAAI/article/view/5541>.
- 2 Jennifer Badham, Frank Kee, and Ruth F. Hunter. Network structure influence on simulated network interventions for behaviour change. *Social Networks*, 64:55–62, 2021.
- 3 Paul Balister, Béla Bollobás, J. Robert Johnson, and Mark Walters. Random majority percolation. *Random Structures & Algorithms*, 36(3):315–340, 2010.
- 4 George Barmpalias, Richard Elwes, and Andrew Lewis-Pye. Tipping points in 1-dimensional Schelling models with switching agents. *Journal of Statistical Physics*, 158(4):1572–9613, 2015.
- 5 George Barmpalias, Richard Elwes, and Andrew Lewis-Pye. Unperturbed Schelling segregation in two or three dimensions. *Journal of Statistical Physics*, 164(6):1460–1487, 2016.

- 6 George Barmpalias, Richard Elwes, and Andrew Lewis-Pye. Minority population in the one-dimensional Schelling model of segregation. *Journal of Statistical Physics*, 173(5):1572–9613, 2018.
- 7 George Barmpalias, Richard Elwes, and Andy Lewis-Pye. Digital morphogenesis via Schelling segregation. In *FOCS'14*, pages 156–165, 2014.
- 8 Stephen Benard and Robb Willer. A wealth and status-based model of residential segregation. *Journal of Mathematical Sociology*, 31(2):149–174, 2007.
- 9 Itzhak Benenson, Erez Hatna, and Ehud Or. From Schelling to spatially explicit modeling of urban ethnic and economic residential dynamics. *Sociological Methods and Research*, 37(4):463–497, 2009.
- 10 Prateek Bhakta, Sarah Miracle, and Dana Randall. Clustering and mixing times for segregation models on \mathbb{Z}^2 . In *SODA'14*, pages 327–340, 2014.
- 11 Davide Bilò, Vittorio Bilò, Pascal Lenzner, and Louise Molitor. Topological influence and locality in swap Schelling games. In *MFCS'20*, pages 15:1–15:15, 2020.
- 12 Thomas Bläsius, Tobias Friedrich, Martin S. Krejca, and Louise Molitor. The flip Schelling process on random geometric and Erdős-Rényi graphs, 2021. [arXiv:2102.09856](https://arxiv.org/abs/2102.09856).
- 13 Christina Brandt, Nicole Immorlica, Gautam Kamath, and Robert Kleinberg. An analysis of one-dimensional Schelling segregation. In *STOC'12*, pages 789–804, 2012.
- 14 Paulo R. A. Campos, Viviane M. de Oliveira, and F. G. Brady Moreira. Small-world effects in the majority-vote model. *Physical Review E*, 67:026104, February 2003.
- 15 Claudio Castellano, Santo Fortunato, and Vittorio Loreto. Statistical physics of social dynamics. *Review of Modern Physics*, 81:591–646, 2009.
- 16 Hau Chan, Mohammad T. Irfan, and Cuong Viet Than. Schelling models with localized social influence: A game-theoretic framework. In *AAMAS'20*, pages 240–248, 2020.
- 17 Ankit Chauhan, Pascal Lenzner, and Louise Molitor. Schelling segregation with strategic agents. In *SAGT'18*, pages 137–149. Springer, 2018.
- 18 William A. V. Clark. Residential segregation in american cities: A review and interpretation. *Population Research and Policy Review*, 5(2):95–127, 1986.
- 19 Vasco Cortez and Sergio Rica. Dynamics of the Schelling social segregation model in networks. *Procedia Computer Science*, 61:60–65, 2015.
- 20 Richard Durrett and Jeffrey E. Steif. Fixation results for threshold voter systems. *Annals of Probability*, 21(1):232–247, 1993.
- 21 Hagen Echezell, Tobias Friedrich, Pascal Lenzner, Louise Molitor, Marcus Pappik, Friedrich Schöne, Fabian Sommer, and David Stangl. Convergence and hardness of strategic Schelling segregation. In *WINE'19*, pages 156–170, 2019.
- 22 Edith Elkind, Jiarui Gan, Atushi Igarashi, Warut Suksompong, and Alexandros A. Voudouris. Schelling games on graphs. In *IJCAI'19*, pages 266–272, 2019.
- 23 Paul Erdős and Alfréd Rényi. On random graphs I. *Publicationes Mathematicae Debrecen*, 6:290–297, 1959.
- 24 William Feller. *An Introduction to Probability Theory and Its Applications*, volume 1. John Wiley & Sons, 3rd edition, 1968.
- 25 Mark A Fossett. Simseg—a computer program to simulate the dynamics of residential segregation by social and ethnic status. *Race and Ethnic Studies Institute Technical Report and Program, Texas A&M University*, 1998.
- 26 Linton C. Freeman. Segregation in social networks. *Sociological Methods & Research*, 6(4):411–429, 1978.
- 27 Laetitia Gauvin, Jean-Pierre Nadal, and Jean Vannimenus. Schelling segregation in an open city: A kinetically constrained blume-emery-griffiths spin-1 system. *Physical Review E*, 81:066120, 2010.
- 28 Stefan Gerhold, Lev Glebsky, Carsten Schneider, Howard Weiss, and Burkhard Zimmermann. Computing the complexity for Schelling segregation models. *Communications in Nonlinear Science and Numerical Simulation*, 13(10):2236–2245, 2008.

- 29 Edgar N. Gilbert. Random graphs. *Annals of Mathematical Statistics*, 30(4):1141–1144, 1959.
- 30 Mark S. Granovetter. Threshold models of collective behavior. *American Journal of Sociology*, 83:1420–1443, 1978.
- 31 Adam Douglas Henry, Paweł Prałat, and Cun-Quan Zhang. Emergence of segregation in evolving social networks. *Proceedings of the National Academy of Sciences*, 108(21):8605–8610, 2011.
- 32 Nicole Immorlica, Robert Kleinberg, Brendan Lucier, and Morteza Zadomighaddam. Exponential segregation in a two-dimensional Schelling model with tolerant individuals. In *SODA'17*, pages 984–993, 2017.
- 33 Panagiotis Kanellopoulos, Maria Kyropoulou, and Alexandros A. Voudouris. Modified Schelling games. In *SAGT'20*, pages 241–256, 2020. doi:10.1007/978-3-030-57980-7_16.
- 34 David Kempe, Jon Kleinberg, and Éva Tardos. Maximizing the spread of influence through a social network. In *Proceedings of the Ninth ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '03, pages 137–146, 2003.
- 35 Thomas M. Liggett. Coexistence in threshold voter models. *The Annals of Probability*, 22(2):764–802, 1994.
- 36 Thomas M. Liggett. *Voter Models*, pages 139–208. Springer, 1999.
- 37 Francisco W.S. Lima, Alexandre O. Sousa, and Muneer A. Sumuor. Majority-vote on directed Erdős–Rényi random graphs. *Physica A: Statistical Mechanics and its Applications*, 387(14):3503–3510, 2008.
- 38 Michael W. Macy. Chains of cooperation: Threshold effects in collective action. *American Sociological Review*, 56(6):730–747, 1991.
- 39 Hamed Omidvar and Massimo Franceschetti. Self-organized segregation on the grid. *Journal of Statistical Physics*, 170(4):1572–9613, 2018.
- 40 Hamed Omidvar and Massimo Franceschetti. Shape of diffusion and size of monochromatic region of a two-dimensional spin system. In *STOC'18*, pages 100–113, 2018.
- 41 Romans Pancs and Nicolaas J. Vriend. Schelling's spatial proximity model of segregation revisited. *Journal of Public Economics*, 91(1):1–24, 2007.
- 42 Mathew Penrose. *Random Geometric Graphs*. Oxford University Press, 1st edition, 2003.
- 43 Luiz F. C. Pereira and F. G. Brady Moreira. Majority-vote model on random graphs. *Physical Review E*, 71:016123, 2005.
- 44 Alexis Poindron. A general model of binary opinions updating. *Mathematical Social Sciences*, 109:52–76, 2021.
- 45 Tim Rogers and Alan J McKane. A unified framework for Schelling's model of segregation. *Journal of Statistical Mechanics: Theory and Experiment*, 2011(07):P07006, 2011.
- 46 Thomas C. Schelling. Models of segregation. *American Economic Review*, 59(2):488–93, 1969.
- 47 Thomas C Schelling. *Micromotives and Macrobehavior*. WW Norton & Company, 2006.
- 48 Sorin Solomon and Dietrich Stauffer. Ising, Schelling and self-organising segregation. *The European Physical Journal B*, 57(4):473–479, 2007.
- 49 Michèle J. White. Segregation and diversity measures in population distribution. *Population index*, 52 2:198–221, 1986.
- 50 Zhi-Xi Wu and Petter Holme. Majority-vote model on hyperbolic lattices. *Physical Review E*, 81:011133, 2010.
- 51 H. Peyton Young. *Individual strategy and social structure: An evolutionary theory of institutions*. Princeton University Press Princeton, N.J, 1998.